

81. Continued

$$(e) v(t) = 3t^2 - 2 > 0$$

$$3t^2 > 2$$

$$t > \frac{\sqrt{6}}{2}$$

$$82. (a) \frac{d}{dx} e^u = e^u \frac{du}{dx} \text{ where } u = x$$

$$\frac{d}{dx} \frac{e^x + e^{-x}}{2} = \frac{e^x - e^{-x}}{2}$$

$$(b) \frac{d}{dx} \frac{e^x - e^{-x}}{2} = \frac{e^x + e^{-x}}{2}$$

$$(c) y(1) = \frac{e^1 + e^{-1}}{2} = 1.543$$

$$y'(1) = \frac{e^1 - e^{-1}}{2} = 1.175$$

$$y = 1.175(x-1) + 1.543$$

$$y = 1.175x + 0.368$$

$$(d) m_2 = -\frac{1}{m_1} = -\frac{1}{1.175} = -0.851$$

$$y = -0.851(x-1) + 1.543$$

$$y = -0.851x + 2.394$$

$$(e) y' = 0 = \frac{e^x - e^{-x}}{2}$$

$$0 = e^x - e^{-x}$$

$$e^x = e^{-x}$$

$$x = -x \text{ or } x = 0$$

$$83. (a) 1 - x^2 > 0$$

$$x^2 > 1, -1 < x < 1$$

$$(b) f'(x) = \frac{d}{dx} \ln(1-x^2) \quad u = 1-x^2$$

$$\frac{d}{dx} \ln(u) = \frac{1}{u} \frac{du}{dx} \quad \frac{du}{dx} = -2x$$

$$= \frac{-2x}{(1-x^2)}$$

$$(c) 1 - x^2 > 0, -1 < x < 1$$

$$(d) y' \left(\frac{1}{2} \right) = \frac{-2 \left(\frac{1}{2} \right)}{1 - \left(\frac{1}{2} \right)^2} = -\frac{1}{3/4} = -4/3$$

Chapter 4

Applications of Derivatives

Section 4.1 Extreme Values of Functions
(pp. 187–195)


Exploration 1 Finding Extreme Values

1. From the graph we can see that there are three critical points: $x = -1, 0, 1$.

Critical point values: $f(-1) = 0.5, f(0) = 0, f(1) = 0.5$


Endpoint values: $f(-2) = 0.4, f(2) = 0.4$

Thus f has absolute maximum value of 0.5 at $x = -1$ and $x = 1$, absolute minimum value of 0 at $x = 0$, and local minimum value of 0.4 at $x = -2$ and $x = 2$.



$[-2, 2]$ by $[-1, 1]$

2. The graph of f' has zeros at $x = -1$ and $x = 1$ where the graph of f has local extreme values. The graph of f' is not defined at $x = 0$, another extreme value of the graph of f .



$[-2, 2]$ by $[-1, 1]$

3. Using the chain rule and $\frac{d}{dx}(|x|) = \frac{|x|}{x}$, we find

$$\frac{df}{dx} = \frac{|x|}{x} \cdot \frac{1-x^2}{(x^2+1)^2}$$

Quick Review 4.1

$$1. f'(x) = \frac{1}{2\sqrt{4-x}} \cdot \frac{d}{dx}(4-x) = \frac{-1}{2\sqrt{4-x}}$$

$$2. f'(x) = \frac{d}{dx} 2(9-x^2)^{-1/2} = -(9-x^2)^{-3/2} \cdot \frac{d}{dx}(9-x^2)$$

$$= -(9-x^2)^{-3/2}(-2x) = \frac{2x}{(9-x^2)^{3/2}}$$

$$3. g'(x) = -\sin(\ln x) \cdot \frac{d}{dx} \ln x = -\frac{\sin(\ln x)}{x}$$

$$4. h'(x) = e^{2x} \cdot \frac{d}{dx} 2x = 2e^{2x}$$

5. Graph (c), since this is the only graph that has positive slope at c .

6. Graph (b), since this is the only graph that represents a differentiable function at a and b and has negative slope at c .

7. Graph (d), since this is the only graph representing a function that is differentiable at b but not at a .

8. Graph (a), since this is the only graph that represents a function that is not differentiable at a or b .

9. As $x \rightarrow 3^-$, $\sqrt{9-x^2} \rightarrow 0^+$. Therefore, $\lim_{x \rightarrow 3^-} f(x) = \infty$.

10. As $x \rightarrow 3^+$, $\sqrt{9-x^2} \rightarrow 0^+$. Therefore, $\lim_{x \rightarrow 3^+} f(x) = \infty$.

$$11. (a) \frac{d}{dx}(x^3 - 2x) = 3x^2 - 2$$

$$f'(1) = 3(1)^2 - 2 = 1$$

$$(b) \frac{d}{dx}(x+2) = 1$$

$$f'(3) = 1$$

(c) Left-hand derivative:

$$\lim_{h \rightarrow 0^-} \frac{f(2+h) - f(2)}{h} = \lim_{h \rightarrow 0^-} \frac{[(2+h)^3 - 2(2+h)] - 4}{h}$$

$$= \lim_{h \rightarrow 0^-} \frac{h^3 + 6h^2 + 10h}{h}$$

$$= \lim_{h \rightarrow 0^-} (h^2 + 6h + 10)$$

$$= 10$$

Right-hand derivative:

$$\lim_{h \rightarrow 0^+} \frac{f(2+h) - f(2)}{h} = \lim_{h \rightarrow 0^+} \frac{[(2+h) + 2] - 4}{h}$$

$$= \lim_{h \rightarrow 0^+} \frac{h}{h}$$

$$= \lim_{h \rightarrow 0^+} 1$$

$$= 1$$

Since the left- and right-hand derivatives are not equal,

$f'(2)$ is undefined.

12. (a) The domain is $x \neq 2$. (See the solution for 11.(c).)

$$(b) f'(x) = \begin{cases} 3x^2 - 2, & x < 2 \\ 1, & x > 2 \end{cases}$$

Section 4.1 Exercises

1. Minima at $(-2, 0)$ and $(2, 0)$, maximum at $(0, 2)$

3. Maximum at $(0, 5)$. Note that there is no minimum since the endpoint $(2, 0)$ is excluded from the graph.

5. Maximum at $x = b$, minimum at $x = c_2$;

The Extreme Value Theorem applies because f is continuous on $[a, b]$, so both the maximum and minimum exist.

7. Maximum at $x = c$, no minimum;

The Extreme Value Theorem does not apply, because the function is not defined on a closed interval.

9. Maximum at $x = c$, minimum at $x = a$;

The Extreme Value Theorem does not apply, because the function is not continuous.

11. The first derivative $f'(x) = -\frac{1}{x^2} + \frac{1}{x}$ has a zero at $x = 1$.

Critical point value: $f(1) = 1 + \ln 1 = 1$

Endpoint values: $f(0.5) = 2 + \ln 0.5 \approx 1.307$

$$f(4) = \frac{1}{4} + \ln 4 \approx 1.636$$

Maximum value is $\frac{1}{4} + \ln 4$ at $x = 4$;

minimum value is 1 at $x = 1$;

local maximum at $\left(\frac{1}{2}, 2 - \ln 2\right)$;

13. The first derivative $h'(x) = \frac{1}{x+1}$ has no zeros, so we need

only consider the endpoints.

$$h(0) = \ln 1 = 0 \quad h(3) = \ln 4$$

Maximum value is $\ln 4$ at $x = 3$;

minimum value is 0 at $x = 0$.

15. The first derivative $f'(x) = \cos\left(x + \frac{\pi}{4}\right)$, has zeros

$$\text{at } x = \frac{\pi}{4}, x = \frac{5\pi}{4}.$$

$$\text{Critical point values: } x = \frac{\pi}{4} \quad f(x) = 1$$

$$x = \frac{5\pi}{4} \quad f(x) = -1$$

$$\text{Endpoint values: } x = 0 \quad f(x) = \frac{1}{\sqrt{2}}$$

$$x = \frac{7\pi}{4} \quad f(x) = 0$$


Maximum value is 1 at $x = \frac{\pi}{4}$;

minimum value is -1 at $x = \frac{5\pi}{4}$;

15. Continued

local minimum at $\left(0, \frac{1}{\sqrt{2}}\right)$;

local maximum at $\left(\frac{7\pi}{4}, 0\right)$



17. The first derivative $f'(x) = \frac{2}{5}x^{-3/5}$ is never zero but is

undefined at $x = 0$.


Critical point value: $x = 0 \quad f(x) = 0$

Endpoint value: $x = -3 \quad f(x) = (-3)^{2/5}$
 $= 3^{2/5} \approx 1.552$


Since $f(x) > 0$ for $x \neq 0$, the critical point at $x = 0$ is a local minimum, and since $f(x) \leq (-3)^{2/5}$ for $-3 \leq x < 1$, the endpoint value at $x = -3$ is a global maximum.

Maximum value is $3^{2/5}$ at $x = -3$;

minimum value is 0 at $x = 0$.





19.




$[-2, 6]$ by $[-2, 4]$

Minimum value is 1 at $x = 2$.



21.




$[-6, 6]$ by $[-5, 20]$

To find the exact values, note that


$y' = 3x^2 + 2x - 8 = (3x - 4)(x + 2)$, which is zero when

$x = -2$ or $x = \frac{4}{3}$. Local maximum at $(-2, 17)$; local minimum

at $\left(\frac{4}{3}, -\frac{41}{27}\right)$





23.



$[-4, 4]$ by $[-2, 4]$


Minimum value is 0 at $x = -1$ and at $x = 1$.





$[-1.5, 1.5]$ by $[-0.5, 3]$



The minimum value is 1 at $x = 0$.



$[-4.7, 4.7]$ by $[-3.1, 3.1]$

Maximum value is 2 at $x = 1$;



minimum value is 0 at $x = -1$ and at $x = 3$.



$[-5, 5]$ by $[-0.7, 0.7]$

Maximum value is $\frac{1}{2}$ at $x = 1$;

minimum value is $-\frac{1}{2}$ at $x = -1$.





$[-6, 6]$ by $[0, 12]$

Maximum value is 11 at $x = 5$;

minimum value is 5 on the interval $[-3, 2]$;



local maximum at $(-5, 9)$



$[-6, 6]$ by $[-6, 6]$

Maximum value is 5 on the interval $[3, \infty)$;

minimum value is -5 on the interval $(-\infty, -2]$.




$[-4, 4]$ by $[-3, 3]$


$$y' = x^{2/3}(1) + \frac{2}{3}x^{-1/3}(x+2) = \frac{5x+4}{3\sqrt[3]{x}}$$

35. Continued

| crit. pt. | derivative | extremum | value |
|--------------------|------------|-----------|---------------------------------------|
| $x = -\frac{4}{5}$ | 0 | local max | $\frac{12}{25}10^{1/3} \approx 1.034$ |
| $x = 0$ | undefined | local min | 0 |




39.




$[-4.7, 4.7]$ by $[0, 6.2]$

$$y' = \begin{cases} -2, & x < 1 \\ 1, & x > 1 \end{cases}$$

| crit. pt. | derivative | extremum | value |
|-----------|------------|----------|-------|
| $x = 1$ | undefined | minimum | 2 |



37.




$[-2.35, 2.35]$ by $[-3.5, 3.5]$


$$y' = x \cdot \frac{1}{2\sqrt{4-x^2}}(-2x) + (1)\sqrt{4-x^2}$$

$$= \frac{-x^2 + (4-x^2)}{\sqrt{4-x^2}} = \frac{4-2x^2}{\sqrt{4-x^2}}$$

| crit. pt. | derivative | extremum | value |
|-----------------|------------|-----------|-------|
| $x = -2$ | undefined | local max | 0 |
| $x = -\sqrt{2}$ | 0 | minimum | -2 |
| $x = \sqrt{2}$ | 0 | maximum | 2 |
| $x = 2$ | undefined | local min | 0 |



41.



$[-4, 6]$ by $[-2, 6]$

$$y' = \begin{cases} -2x - 2, & x < 1 \\ -2x + 6, & x > 1 \end{cases}$$

| crit. pt. | derivative | extremum | value |
|-----------|------------|-----------|-------|
| $x = -1$ | 0 | maximum | 5 |
| $x = 1$ | undefined | local min | 1 |
| $x = 3$ | 0 | maximum | 5 |

- (b) The largest possible volume of the box is 144 cubic units, and it occurs when $x = 2$.

45. False. For example, the maximum could occur at a corner, where $f'(c)$ would not exist.

$$47. E. \frac{d}{dx}(4x - x^2 + 6) = 4 - 2x$$

$$4 - 2x = 0$$

$$x = 2$$

$$f(2) = 4(2) - (2)^2 + 6 = 10$$

$$49. B. \frac{d}{dx}(x^3 - 6x + 5) = 3x^2 - 6$$

$$3x^2 - 6 = 0$$

$$x = \pm\sqrt{2}$$

51. (a) No, since $f'(x) = \frac{2}{3}(x-2)^{-1/3}$, which is undefined at $x = 2$.

- (b) The derivative is defined and nonzero for all $x \neq 2$. Also, $f(2) = 0$ and $f(x) > 0$ for all $x \neq 2$.

- (c) No, $f(x)$ need not have a global maximum because its domain is all real numbers. Any restriction of f to a closed interval of the form $[a, b]$ would have both a maximum value and a minimum value on the interval.

- (d) The answers are the same as (a) and (b) with 2 replaced by a .

43. (a) $V(x) = 160x - 52x^2 + 4x^3$


$$V'(x) = 160 - 104x + 12x^2 = 4(x-2)(3x-20)$$

The only critical point in the interval $(0, 5)$ is at $x = 2$.

The maximum value of $V(x)$ is 144 at $x = 2$.


53. (a) $f'(x) = 3ax^2 + 2bx + c$ is a quadratic, so it can have 0, 1, or 2 zeros, which would be the critical points of f .

Examples:




$[-3, 3]$ by $[-5, 5]$

The function $f(x) = x^3 - 3x$ has two critical points at $x = -1$ and $x = 1$.



$[-3, 3]$ by $[-5, 5]$

The function $f(x) = x^3 - 1$ has one critical point at $x = 0$.




$[-3, 3]$ by $[-5, 5]$

The function $f(x) = x^3 + x$ has no critical points.

- (b) The function can have either two local extreme values or no extreme values. (If there is only one critical point, the cubic function has no extreme values.)

55. (a)



$[-0.1, 0.6]$ by $[-1.5, 1.5]$

$f(0) = 0$ is not a local extreme value because in any open interval containing $x = 0$, there are infinitely many points where $f(x) = 1$ and where $f(x) = -1$.

- (b) One possible answer, on the interval $[0, 1]$:

$$f(x) = \begin{cases} (1-x) \cos \frac{1}{1-x}, & 0 \leq x < 1 \\ 0, & x = 1 \end{cases}$$

This function has no local extreme value at $x = 1$. Note that it is continuous on $[0, 1]$.

Section 4.2 Mean Value Theorem (pp. 196–204)

Quick Review 4.2

1. $2x^2 - 6 < 0$

$$2x^2 < 6$$

$$x^2 < 3$$

$$-\sqrt{3} < x < \sqrt{3}$$

Interval: $(-\sqrt{3}, \sqrt{3})$

2. $3x^2 - 6 > 0$

$$3x^2 > 6$$

$$x^2 > 2$$

$$x < -\sqrt{2} \text{ or } x > \sqrt{2}$$

Intervals: $(-\infty, -\sqrt{2}) \cup (\sqrt{2}, \infty)$

3. Domain: $8 - 2x^2 \geq 0$

$$8 \geq 2x^2$$

$$4 \geq x^2$$

$$-2 \leq x \leq 2$$

The domain is $[-2, 2]$.

4. f is continuous for all x in the domain, or, in the interval $[-2, 2]$.

5. f is differentiable for all x in the interior of its domain, or, in the interval $(-2, 2)$.
6. We require $x^2 - 1 \neq 0$, so the domain is $x \neq \pm 1$.
7. f is continuous for all x in the domain, or, for all $x \neq \pm 1$.
8. f is differentiable for all x in the domain, or, for all $x \neq \pm 1$.
9. $7 = -2(-2) + C$
 $7 = 4 + C$
 $C = 3$
10. $-1 = (1)^2 + 2(1) + C$
 $-1 = 3 + C$
 $C = -4$

Section 4.2 Exercises

1. (a) Yes.

(b) $f'(x) = \frac{d}{dx} x^2 + 2x - 1 = 2x + 2$
 $2c + 2 = \frac{2 - (-1)}{1 - 0} = 3$
 $c = \frac{1}{2}$.

2. (a) Yes.

(b) $f'(x) = \frac{d}{dx} x^{2/3} = \frac{2}{3} x^{-1/3}$
 $\frac{2}{3} c^{-1/3} = \frac{1 - 0}{1 - 0} = 1$
 $c = \frac{8}{27}$.

3. (a) No. There is a verticle tangent at $x = 0$.4. (a) No. There is a corner at $x = 1$.

5. (a) Yes.

(b) $f'(x) = \frac{d}{dx} \sin^{-1} x = \frac{1}{\sqrt{1-x^2}}$
 $\frac{1}{\sqrt{1-c^2}} = \frac{(\pi/2) - (-\pi/2)}{1 - (-1)} = \frac{\pi}{2}$
 $\sqrt{1-c^2} = \frac{2}{\pi}$
 $c = \sqrt{1 - 4/\pi^2} \approx 0.771$.

6. (a) Yes.

(b) $f'(x) = \frac{d}{dx} \ln(x-1) = \frac{1}{x-1}$
 $\frac{1}{c-1} = \frac{\ln 3 - \ln 1}{4-2}$
 $c = \frac{4-2}{\ln 3 - \ln 1} + 1 \approx 2.820$

7. (a) No. The function is discontinuous at $x = \frac{\pi}{2}$ 8. (a) No. The split function is discontinuous at $x = 1$ 9. (a) The secant line passes through $(0.5, f(0.5)) = (0.5, 2.5)$ and $(2, f(2)) = (2, 2.5)$, so its equation is $y = 2.5$.(b) The slope of the secant line is 0, so we need to find c such that $f'(c) = 0$.

$$1 - c^{-2} = 0$$

$$c^{-2} = 1$$

$$c = 1$$

$$f(c) = f(1) = 2$$

The tangent line has slope 0 and passes through $(1, 2)$, so its equation is $y = 2$.10. (a) The secant line passes through $(1, f(1)) = (1, 0)$ and $(3, f(3)) = (3, \sqrt{2})$, so its slope is

$$\frac{\sqrt{2} - 0}{3 - 1} = \frac{\sqrt{2}}{2} = \frac{1}{\sqrt{2}}$$

The equation is $y = \frac{1}{\sqrt{2}}(x-1) + 0$ or $y = \frac{1}{\sqrt{2}}x - \frac{1}{\sqrt{2}}$, or $y \approx 0.707x - 0.707$.(b) We need to find c such that $f'(c) = \frac{1}{\sqrt{2}}$.

$$\frac{1}{2\sqrt{c-1}} = \frac{1}{\sqrt{2}}$$

$$2\sqrt{c-1} = \sqrt{2}$$

$$c-1 = \frac{1}{2}$$

$$c = \frac{3}{2}$$

$$f(c) = f\left(\frac{3}{2}\right) = \sqrt{\frac{1}{2}} = \frac{1}{\sqrt{2}}$$

The tangent line has slope $\frac{1}{\sqrt{2}}$ and passes through $\left(\frac{3}{2}, \frac{1}{\sqrt{2}}\right)$. Its equation is $y = \frac{1}{\sqrt{2}}\left(x - \frac{3}{2}\right) + \frac{1}{\sqrt{2}}$ or $y = \frac{1}{\sqrt{2}}x - \frac{1}{2\sqrt{2}}$, or $y \approx 0.707x - 0.354$.

11. Because the trucker's average speed was 79.5 mph, and by then Mean Value Theorem, the trucker must have been going that speed at least once during the trip.

12. Let $f(t)$ denote the temperature indicated after t seconds. We assume that $f'(t)$ is defined and continuous for $0 \leq t \leq 20$. The average rate of change is $10.6^\circ\text{F}/\text{sec}$. Therefore, by the Mean Value Theorem, $f'(c) = 10.6^\circ\text{F}/\text{sec}$ for some value of c in $[0, 20]$. Since the temperature was constant before $t = 0$, we also know that $f'(0) = 0^\circ\text{F}/\text{min}$. But f' is continuous, so by the Intermediate Value Theorem, the rate of change $f'(t)$ must have been $10.1^\circ\text{F}/\text{sec}$ at some moment during the interval.

13. Because its average speed was approximately 7.667 knots, and by the Mean Value Theorem, it must have been going that speed at least once during the trip.

14. The runner's average speed for the marathon was approximately 11.909 mph. Therefore, by the Mean Value Theorem, the runner must have been going that speed at least once during the marathon. Since the initial speed and final speed are both 0 mph and the runner's speed is continuous, by the Intermediate Value Theorem, the runner's speed must have been 11 mph at least twice.

15. (a) $f'(x) = 5 - 2x$

Since $f'(x) > 0$ on $\left(-\infty, \frac{5}{2}\right)$, $f'(x) = 0$ at $x = \frac{5}{2}$, and

$f'(x) < 0$ on $\left(\frac{5}{2}, \infty\right)$, we know that $f(x)$ has a local

maximum at $x = \frac{5}{2}$. Since $f\left(\frac{5}{2}\right) = \frac{25}{4}$, the local

maximum occurs at the point $\left(\frac{5}{2}, \frac{25}{4}\right)$. (This is also a global maximum.)

(b) Since $f'(x) > 0$ on $\left(-\infty, \frac{5}{2}\right)$, $f(x)$ is increasing on

$\left(-\infty, \frac{5}{2}\right]$.

(c) Since $f'(x) < 0$ on $\left(\frac{5}{2}, \infty\right)$, $f(x)$ is decreasing on

$\left[\frac{5}{2}, \infty\right)$.

16. (a) $g'(x) = 2x - 1$

Since $g'(x) < 0$ on $\left(-\infty, \frac{1}{2}\right)$, $g'(x) = 0$ at $x = \frac{1}{2}$, and

$g'(x) > 0$ on $\left(\frac{1}{2}, \infty\right)$, we know that $g(x)$ has a local

minimum at $x = \frac{1}{2}$.

Since $g\left(\frac{1}{2}\right) = -\frac{49}{4}$, the local minimum occurs at the

point $\left(\frac{1}{2}, -\frac{49}{4}\right)$. (This is also a global minimum.)

(b) Since $g'(x) > 0$ on $\left(\frac{1}{2}, \infty\right)$, $g(x)$ is increasing on

$\left[\frac{1}{2}, \infty\right)$.

(c) Since $g'(x) < 0$ on $\left(-\infty, \frac{1}{2}\right)$, $g(x)$ is decreasing on

$\left(-\infty, \frac{1}{2}\right]$.

17. (a) $h'(x) = -\frac{2}{x^2}$

Since $h'(x)$ is never zero or undefined only where $h(x)$ is undefined, there are no critical points. Also, the domain $(-\infty, 0) \cup (0, \infty)$ has no endpoints. Therefore, $h(x)$ has no local extrema.

(b) Since $h'(x)$ is never positive, $h(x)$ is not increasing on any interval.

(c) Since $h'(x) < 0$ on $(-\infty, 0) \cup (0, \infty)$, $h(x)$ is decreasing on $(-\infty, 0)$ and on $(0, \infty)$.

18. (a) $k'(x) = -\frac{2}{x^3}$

Since $k'(x)$ is never zero and is undefined only where $k(x)$ is undefined, there are no critical points. Also, the domain $(-\infty, 0) \cup (0, \infty)$ has no endpoints. Therefore, $k(x)$ has no local extrema.

(b) Since $k'(x) > 0$ on $(-\infty, 0)$, $k(x)$ is increasing on $(-\infty, 0)$.

(c) Since $k'(x) < 0$ on $(0, \infty)$, $k(x)$ is decreasing on $(0, \infty)$.

19. (a) $f'(x) = 2e^{2x}$

Since $f'(x)$ is never zero or undefined, and the domain of $f(x)$ has no endpoints, $f(x)$ has no extrema.

(b) Since $f'(x)$ is always positive, $f(x)$ is increasing on $(-\infty, \infty)$.

(c) Since $f'(x)$ is never negative, $f(x)$ is not decreasing on any interval.

20. (a) $f'(x) = -0.5e^{-0.5x}$

Since $f'(x)$ is never zero or undefined, and the domain of $f(x)$ has no endpoints, $f(x)$ has no extrema.

(b) Since $f'(x)$ is never positive, $f(x)$ is not increasing on any interval.

(c) Since $f'(x)$ is always negative, $f(x)$ is decreasing on $(-\infty, \infty)$.

21. (a) $y' = -\frac{1}{2\sqrt{x+2}}$

In the domain $[-2, \infty)$, y' is never zero and is undefined only at the endpoint $x = -2$. The function y has a local maximum at $(-2, 4)$. (This is also a global maximum.)

(b) Since y' is never positive, y is not increasing on any interval.


(c) Since y' is negative on $(-2, \infty)$, y is decreasing on $[-2, \infty)$.

22. (a) $y' = 4x^3 - 20x = 4x(x + \sqrt{5})(x - \sqrt{5})$

The function has critical points at $x = -\sqrt{5}$, $x = 0$, and $x = \sqrt{5}$. Since $y' < 0$ on $(-\infty, -\sqrt{5})$ and $(0, \sqrt{5})$ and $y' > 0$ on $(-\sqrt{5}, 0)$ and $(\sqrt{5}, \infty)$, the points at $x = \pm\sqrt{5}$ are local minima and the point at $x = 0$ is a local maximum. Thus, the function has a local maximum at $(0, 9)$ and local minima at $(-\sqrt{5}, -16)$ and $(\sqrt{5}, -16)$. (These are also global minima.)

(b) Since $y' > 0$ on $(-\sqrt{5}, 0)$ and $(\sqrt{5}, \infty)$, y is increasing on $[-\sqrt{5}, 0]$ and $[\sqrt{5}, \infty)$.

(c) Since $y' < 0$ on $(-\infty, -\sqrt{5})$ and $(0, \sqrt{5})$, y is decreasing on $(-\infty, -\sqrt{5}]$ and $[0, \sqrt{5}]$.



$[-4.7, 4.7]$ by $[-3.1, 3.1]$


(a)
$$f'(x) = x \cdot \frac{1}{2\sqrt{4-x}} (-1) + \sqrt{4-x}$$

$$= \frac{-3x+8}{2\sqrt{4-x}}$$

The local extrema occur at the critical point $x = \frac{8}{3}$ and at the endpoint $x = 4$. There is a local (and absolute) maximum at $(\frac{8}{3}, \frac{16}{3\sqrt{3}})$ or approximately $(2.67, 3.08)$, and a local minimum at $(4, 0)$.

(b) Since $f'(x) > 0$ on $(-\infty, \frac{8}{3})$, $f(x)$ is decreasing on $(-\infty, \frac{8}{3}]$.

(c) Since $f'(x) < 0$ on $(\frac{8}{3}, 4)$, $f(x)$ is decreasing on $[\frac{8}{3}, 4]$.




$[-5, 5]$ by $[-15, 15]$

(a)
$$g'(x) = x^{1/3}(1) + \frac{1}{3}x^{-2/3}(x+8) = \frac{4x+8}{3x^{2/3}}$$

The local extrema can occur at the critical points $x = -2$ and $x = 0$, but the graph shows that no extrema occurs at $x = 0$. There is a local (and absolute) minimum at $(-2, -6\sqrt[3]{2})$ or approximately $(-2, -7.56)$.

(b) Since $g'(x) > 0$ on the intervals $(-2, 0)$ and $(0, \infty)$, and $g(x)$ is continuous at $x = 0$, $g(x)$ is increasing on $[-2, \infty)$.

(c) Since $g'(x) < 0$ on the interval $(-\infty, -2)$, $g(x)$ is decreasing on $(-\infty, -2]$.



$[-5, 5]$ by $[-0.4, 0.4]$

(a)
$$h'(x) = \frac{(x^2+4)(-1) - (-x)(2x)}{(x^2+4)^2} = \frac{x^2-4}{(x^2+4)^2}$$

$$= \frac{(x+2)(x-2)}{(x^2+4)^2}$$


The local extrema occur at the critical points, $x = \pm 2$.

There is a local (and absolute) maximum at $(-2, \frac{1}{4})$

and a local (and absolute) minimum at $(2, -\frac{1}{4})$.

(b) Since $h'(x) > 0$ on $(-\infty, -2)$ and $(2, \infty)$, $h(x)$ is increasing on $(-\infty, -2]$ and $[2, \infty)$.

(c) Since $h'(x) < 0$ on $(-2, 2)$, $h(x)$ is decreasing on $[-2, 2]$.




$[-4.7, 4.7]$ by $[-3.1, 3.1]$

(a)
$$k'(x) = \frac{(x^2-4)(1) - x(2x)}{(x^2-4)^2} = -\frac{x^2+4}{(x^2-4)^2}$$

Since $k'(x)$ is never zero and is undefined only where $k(x)$ is undefined, there are no critical points. Since there are no critical points and the domain includes no endpoints, $k(x)$ has no local extrema.

(b) Since $k'(x)$ is never positive, $k(x)$ is not increasing on any interval.

(c) Since $k'(x)$ is negative wherever it is defined, $k(x)$ is decreasing on each interval of its domain; on $(-\infty, -2)$, $(-2, 2)$, and $(2, \infty)$.




(a) $f'(x) = 3x^2 - 2 + 2\sin x$

Note that $3x^2 - 2 > 2$ for $|x| \geq 1.2$ and $|2\sin x| \leq 2$ for all x , so $f'(x) > 0$ for $|x| \geq 1.2$. Therefore, all critical points occur in the interval $(-1.2, 1.2)$, as suggested by the graph. Using grapher techniques, there is a local maximum at approximately $(-1.126, -0.036)$, and a local minimum at approximately $(0.559, -2.639)$.

(b) $f(x)$ is increasing on the intervals $(-\infty, -1.126]$ and $[0.559, \infty)$, where the interval endpoints are approximate.

(c) $f(x)$ is decreasing on the interval $[-1.126, 0.559]$, where the interval endpoints are approximate.



(a) $g'(x) = 2 - \sin x$

Since $1 \leq g'(x) \leq 3$ for all x , there are no critical points. Since there are no critical points and the domain has no endpoints, there are no local extrema.

(b) Since $g'(x) > 0$ for all x , $g(x)$ is increasing on $(-\infty, \infty)$.

(c) Since $g'(x)$ is never negative, $g(x)$ is not decreasing on any interval.

29. $f(x) = \frac{x^2}{2} + C$

30. $f(x) = 2x + C$

31. $f(x) = x^3 - x^2 + x + C$

32. $f(x) = -\cos x + C$

33. $f(x) = e^x + C$

34. $f(x) = \ln(x-1) + C$

35. $f(x) = \frac{1}{x} + C, x > 0$

$$f(2) = 1$$

$$\frac{1}{2} + C = 1$$

$$C = \frac{1}{2}$$




$$f(x) = \frac{1}{x} + \frac{1}{2}, x > 0$$

36. $f(x) = x^{1/4} + C$
 $f(1) = -2$
 $1^{1/4} + C = -2$
 $1 + C = -2$
 $C = -3$
 $f(x) = x^{1/4} - 3$



37. $f(x) = \ln(x+2) + C$
 $f(-1) = 3$
 $\ln(-1+2) + C = 3$
 $0 + C = 3$
 $C = 3$
 $f(x) = \ln(x+2) + 3$

38. $f(x) = x^2 + x - \sin x + C$
 $f(0) = 3$
 $0 + C = 3$
 $C = 3$
 $f(x) = x^2 + x - \sin x + 3$


39. Possible answers:



40. Possible answers:



40. Continued




[-1, 5] by [-1, 8]




[-1, 5] by [-1, 8]

41. One possible answer:



[-3, 3] by [-15, 15]

42. One possible answer:



[-3, 3] by [-70, 70]

43. (a) Since $v'(t) = 1.6$, $v(t) = 1.6t + C$. But $v(0) = 0$, so $C = 0$ and $v(t) = 1.6t$. Therefore, $v(30) = 1.6(30) = 48$. The rock will be going 48 m/sec.

(b) Let $s(t)$ represent position.

Since $s'(t) = v(t) = 1.6t$, $s(t) = 0.8t^2 + D$. But $s(0) = 0$, so $D = 0$ and $s(t) = 0.8t^2$. Therefore, $s(30) = 0.8(30)^2 = 720$. The rock travels 720 meters in the 30 seconds it takes to hit bottom, so the bottom of the crevasse is 720 meters below the point of release.

(c) The velocity is now given by $v(t) = 1.6t + C$, where $v(0) = 4$. (Note that the sign of the initial velocity is the same as the sign used for the acceleration, since both act in a downward direction.) Therefore, $v(t) = 1.6t + 4$, and $s(t) = 0.8t^2 + 4t + D$, where $s(0) = 0$ and so $D = 0$. Using $s(t) = 0.8t^2 + 4t$ and the known crevasse depth of 720 meters, we solve $s(t) = 720$ to obtain the positive solution $t \approx 27.604$, and so $v(t) = v(27.604) = 1.6(27.604) + 4 \approx 48.166$. The rock will hit bottom after about 27.604 seconds, and it will be going about 48.166 m/sec.

44. (a) We assume the diving board is located at $s = 0$ and the water at $s = 0$, so that downward velocities are positive. The acceleration due to gravity is 9.8 m/sec^2 , so

$v'(t) = 9.8$ and $v(t) = 9.8t + C$. Since $v(0) = 0$, we have $v(t) = 9.8t$. Then the position is given by $s(t)$ where $s'(t) = v(t) = 9.8t$, so $s(t) = 4.9t^2 + D$. Since $s(0) = 0$, we have $s(t) = 4.9t^2$. Solving $s(t) = 10$ gives

$$t^2 = \frac{10}{4.9} = \frac{100}{49}, \text{ so the positive solution is } t = \frac{10}{7}. \text{ The}$$

velocity at this time is $v\left(\frac{10}{7}\right) = 9.8\left(\frac{10}{7}\right) = 14 \text{ m/sec}$.

(b) Again $v(t) = 9.8t + C$, but this time $v(0) = -2$ and so $v(t) = 9.8t - 2$. The $s'(t) = 9.8t - 2$, so $s(t) =$

$$4.9t^2 - 2t + D. \text{ Since } s(0) = 0, \text{ we have } s(t) =$$

$4.9t^2 - 2t$. Solving $s(t) = 10$ gives the positive solution

$$t = \frac{2 + 10\sqrt{2}}{9.8} \approx 1.647 \text{ sec}.$$

The velocity at this time is

$$v\left(\frac{2 + 10\sqrt{2}}{9.8}\right) = 9.8\left(\frac{2 + 10\sqrt{2}}{9.8}\right) - 2 = 10\sqrt{2} \text{ m/sec or}$$

about 14.142 m/sec.

45. Because the function is not continuous on $[0, 1]$. The function does not satisfy the hypotheses of the Mean Value Theorem, and so it need not satisfy the conclusion of the Mean Value Theorem.


46. Because the Mean Value Theorem applies to the function $y = \sin x$ on any interval, and $y = \cos x$ is the derivative of $\sin x$. So, between any two zeros of $\sin x$, its derivative, $\cos x$, must be zero at least once.

47. $f(x)$ must be zero at least once between a and b by the Intermediate Value Theorem. Now suppose that $f(x)$ is zero twice between a and b . Then by the Mean Value Theorem, $f'(x)$ would have to be zero at least once between the two zeros of $f(x)$, but this can't be true since we are given that $f'(x) \neq 0$ on this interval. Therefore, $f(x)$ is zero once and only once between a and b .

48. Let $f(x) = x^4 + 3x + 1$. Then $f(x)$ is continuous and differentiable everywhere. $f'(x) = 4x^3 + 3$, which is never zero between $x = -2$ and $x = -1$. Since $f(-2) = 11$ and $f(-1) = -1$, exercise 47 applies, and $f(x)$ has exactly one zero between $x = -2$ and $x = -1$.

49. Let $f(x) = x + \ln(x + 1)$. Then $f(x)$ is continuous and differentiable everywhere on $[0, 3]$. $f'(x) = 1 + \frac{1}{x+1}$, which is never zero on $[0, 3]$. Now $f(0) = 0$, so $x = 0$ is one solution of the equation. If there were a second solution, $f(x)$ would be zero twice in $[0, 3]$, and by the Mean Value Theorem, $f'(x)$ would have to be zero somewhere between the two zeros of $f(x)$. But this can't happen, since $f'(x)$ is never zero on $[0, 3]$. Therefore, $f(x) = 0$ has exactly one solution in the interval $[0, 3]$.

50. Consider the function $k(x) = f(x) - g(x)$. $k(x)$ is continuous and differentiable on $[a, b]$, and since $k(a) = f(a) - g(a) = 0$ and $k(b) = f(b) - g(b) = 0$, by the Mean Value Theorem, there must be a point c in (a, b) where $k'(c) = 0$. But since $k'(c) = f'(c) - g'(c)$, this means that $f'(c) = g'(c)$, and c is a point where the graphs of f and g have parallel or identical tangent lines.



$(-1, 1)$ by $[-2, 2]$

51. False. For example, the function x^3 is increasing on $(-1, 1)$, but $f'(0) = 0$.
52. True. In fact, f is the increasing on $[a, b]$ by Corollary to the Mean Value Theorem.

53. A. $f'(x) = \frac{\frac{1}{2} - 1}{\frac{\pi}{3}} = -\frac{3}{2\pi}$.


54. B. $f'(x) = \frac{f(4) - f(0)}{4 - 0}$
 $= \frac{3.78 - 2980.96}{4 - 0}$
 $= -744.30$, negative slope.

55. E. $\frac{d}{dx}(2\sqrt{x} - 10)$
 $= \frac{2}{2\sqrt{x}} = \frac{1}{\sqrt{x}}$.

56. D. $x^{3/5}$ is not differentiable at $x = 0$.

57. (a) Increasing: $[-2, -1.3]$ and $[1.3, 2]$;
 decreasing: $[-1.3, 1.3]$;
 local max: $x \approx -1.3$
 local min: $x \approx 1.3$

- (b) Regression equation: $y = 3x^2 - 5$



$[-2.5, 2.5]$ by $[-8, 10]$


- (c) Since $f'(x) = 3x^2 - 5$, we have $f(x) = x^3 - 5x + C$.

But $f(0) = 0$, so $C = 0$. Then $f(x) = x^3 - 5x$.

58. (a) Toward: $0 < t < 2$ and $5 < t < 8$; away: $2 < t < 5$
- (b) A local extremum in this problem is a time/place where Priya changes the direction of her motion.

- (c) Regression equation:

$$y = -0.0820x^3 + 0.9163x^2 - 2.5126x + 3.3779$$



$[-0.5, 8.5]$ by $[-0.5, 5]$

- (d) Using the unrounded values from the regression equation, we obtain

$f'(t) = -0.2459t^2 + 1.8324t - 2.5126$. According to the regression equation, Priya is moving toward the motion detector when $f'(t) < 0$ ($0 < t < 1.81$ and $5.64 < t < 8$), and away from the detector when $f'(t) > 0$ ($1.81 < t < 5.64$).

59. $\frac{f(b) - f(a)}{b - a} = \frac{\frac{1}{b} - \frac{1}{a}}{b - a} = -\frac{1}{ab}$

$f'(c) = -\frac{1}{c^2}$, so $-\frac{1}{c^2} = -\frac{1}{ab}$ and $c^2 = ab$.

Thus, $c = \sqrt{ab}$.

60. $\frac{f(b) - f(a)}{b - a} = \frac{b^2 - a^2}{b - a} = b + a$

$f'(c) = 2c$, so $2c = b + a$ and $c = \frac{a + b}{2}$.

61. By the Mean Value Theorem, $\sin b - \sin a = (\cos c)(b - a)$ for some c between a and b . Taking the absolute value of both sides and using $|\cos c| \leq 1$ gives the result.

62. Apply the Mean Value Theorem to f on $[a, b]$.

Since $f(b) < f(a)$, $\frac{f(b) - f(a)}{b - a}$ is negative, and

hence $f'(x)$ must be negative at some point between a and b .

63. Let $f(x)$ be a monotonic function defined on an interval D . For any two values in D , we may let x_1 be the smaller value and let x_2 be the larger value, so $x_1 < x_2$. Then either $f(x_1) < f(x_2)$ (if f is increasing), or $f(x_1) > f(x_2)$ (if f is decreasing), which means $f(x_1) \neq f(x_2)$. Therefore, f is one-to-one.

Section 4.3 Connecting f' and f'' with the Graph of f (pp. 205–218)


Exploration 1 Finding f from f''

- Any function $f(x) = x^4 - 4x^3 + C$ where C is a real number. For example, let $C = 0, 1, 2$. Their graphs are all vertical shifts of each other.
- Their behavior is the same as the behavior of the function f of Example 8.

Exploration 2 Finding f from f' and f''

1. f has an absolute maximum at $x = 0$ and an absolute minimum of 1 at $x = 4$. We are not given enough information to determine $f(0)$.

2. f has a point of inflection at $x = 2$.



$[-3, 5]$ by $[-5, 20]$

Quick Review 4.3

1. $x^2 - 9 < 0$
 $(x+3)(x-3) < 0$

| Intervals | $x < -3$ | $-3 < x < 3$ | $3 < x$ |
|----------------------|----------|--------------|---------|
| Sign of $(x+3)(x-3)$ | + | - | + |

Solution set: $(-3, 3)$

2. $x^3 - 4x > 0$
 $x(x+2)(x-2) > 0$

| Intervals | $x < -2$ | $-2 < x < 0$ | $0 < x < 2$ | $2 < x$ |
|-----------------------|----------|--------------|-------------|---------|
| Sign of $x(x+2)(x-2)$ | - | + | - | + |

Solution set: $(-2, 0) \cup (2, \infty)$

3. f : all reals

f' : all reals, since $f'(x) = xe^x + e^x$

4. f : all reals

f' : $x \neq 0$, since $f'(x) = \frac{3}{5}x^{-2/5}$

5. f : $x \neq 2$

f' : $x \neq 2$, since $f'(x) = \frac{(x-2)(1)-(x)(1)}{(x-2)^2} = \frac{-2}{(x-2)^2}$

6. f : all reals

f' : $x \neq 0$, since $f'(x) = \frac{2}{5}x^{-3/5}$

7. Left end behavior model: 0

Right end behavior model: $-x^2e^x$

Horizontal asymptote: $y = 0$

8. Left end behavior model: x^2e^{-x}

Right end behavior model: 0

Horizontal asymptote: $y = 0$

9. Left end behavior model: 0

Right end behavior model: 200

Horizontal asymptote: $y = 0$, $y = 200$

10. Left end behavior model: 0

Right end behavior model: 375


Horizontal asymptotes: $y = 0$, $y = 375$

Section 4.3 Exercises

1. $y' = 2x - 1$

| Intervals | $x < \frac{1}{2}$ | $x > \frac{1}{2}$ |
|-----------------|-------------------|-------------------|
| Sign of y' | - | + |
| Behavior of y | Decreasing | Increasing |

Graphical support:




$[-4, 4]$ by $[-3, 3]$

Local (and absolute) minimum at $(\frac{1}{2}, -\frac{5}{4})$

2. $y' = -6x^2 + 12x = -6x(x-2)$

| Intervals | $x < 0$ | $0 < x < 2$ | $2 < x$ |
|-----------------|------------|-------------|------------|
| Sign of y' | - | + | - |
| Behavior of y | Decreasing | Increasing | Decreasing |

Graphical support:



$[-4, 4]$ by $[-6, 6]$


Local maximum: $(2, 5)$;

local minimum: $(0, -3)$

3. $y' = 8x^3 - 8x = 8x(x-1)(x+1)$

| Intervals | $x < -1$ | $-1 < x < 0$ | $0 < x < 1$ | $1 < x$ |
|-----------------|------------|--------------|-------------|------------|
| Sign of y' | - | + | - | + |
| Behavior of y | Decreasing | Increasing | Decreasing | Increasing |

Graphical support:



$[-4, 4]$ by $[-3, 3]$


Local maximum: $(0, 1)$;

local (and absolute) minima: $(-1, -1)$ and $(1, -1)$

4. $y' = xe^{1/x}(-x^{-2}) + e^{1/x} = e^{1/x} \left(1 - \frac{1}{x}\right)$

| | | | |
|-----------------|------------|-------------|------------|
| Intervals | $x < 0$ | $0 < x < 1$ | $1 < x$ |
| Sign of y' | + | - | + |
| Behavior of y | Increasing | Decreasing | Increasing |

Graphical support:




$[-8, 8]$ by $[-6, 6]$

Local minimum: $(1, e)$

5. $y' = x \frac{1}{2\sqrt{8-x^2}}(-2x) + (\sqrt{8-x^2})(1) = \frac{8-2x^2}{\sqrt{8-x^2}}$

| | | | |
|-----------------|----------------------|--------------|--------------------|
| Intervals | $-\sqrt{8} < x < -2$ | $-2 < x < 2$ | $2 < x < \sqrt{8}$ |
| Sign of y' | - | + | - |
| Behavior of y | Decreasing | Increasing | Decreasing |

Graphical support:



$[-3.02, 3.02]$ by $[-6.5, 6.5]$

Local maxima: $(-\sqrt{8}, 0)$ and $(2, 4)$;


local minima: $(-2, -4)$ and $(\sqrt{8}, 0)$

Note that the local extrema at $x = \pm 2$ are also absolute extrema.

6. $y' = \begin{cases} -2x, & x < 0 \\ 2x, & x > 0 \end{cases}$

| | | |
|-----------------|------------|------------|
| Intervals | $x < 0$ | $x > 0$ |
| Sign of y' | + | + |
| Behavior of y | Increasing | Increasing |

Graphical support:



$[-4, 4]$ by $[-3, 6]$

Local minimum: $(0, 1)$


7. $y' = 12x^2 + 42x + 36 = 6(x+2)(2x+3)$

| | | | |
|-----------------|------------|-------------------------|--------------------|
| Intervals | $x < -2$ | $-2 < x < -\frac{3}{2}$ | $-\frac{3}{2} < x$ |
| Sign of y' | + | - | + |
| Behavior of y | Increasing | Decreasing | Increasing |

$y'' = 24x + 42 = 6(4x + 7)$

| | | |
|-----------------|--------------------|--------------------|
| Intervals | $x < -\frac{7}{4}$ | $-\frac{7}{4} < x$ |
| Sign of y'' | - | + |
| Behavior of y | Concave down | Concave up |

Graphical support:



$[-4, 4]$ by $[-80, 20]$

(a) $\left(-\frac{7}{4}, \infty\right)$

(b) $\left(-\infty, -\frac{7}{4}\right)$

8. $y' = -4x^3 + 12x^2 - 4$


Using grapher techniques, the zeros of y' are $x \approx -0.53$, $x \approx 0.65$, and $x \approx 2.88$.

| | | | | |
|-----------------|-------------|--------------------|-------------------|------------|
| Intervals | $x < -0.53$ | $-0.53 < x < 0.65$ | $0.65 < x < 2.88$ | $2.88 < x$ |
| Sign of y' | + | - | + | - |
| Behavior of y | Increasing | Decreasing | Increasing | Decreasing |

$y'' = -12x^2 + 24x = -12x(x-2)$

| | | | |
|-----------------|--------------|-------------|--------------|
| Intervals | $x < 0$ | $0 < x < 2$ | $2 < x$ |
| Sign of y'' | - | + | - |
| Behavior of y | Concave down | Concave up | Concave down |

Graphical support:



$[-2, 4]$ by $[-20, 20]$

(a) $(-\infty, -0.53]$ and $[0.65, 2.88]$

(b) $[-0.53, 0.65]$ and $[2.88, \infty)$

(c) $(0, 2)$

(d) $(-\infty, 0)$ and $(2, \infty)$

8. Continued

(e) Local maxima: $(-0.53, 2.45)$ and $(2.88, 16.23)$; local minimum: $(0.65, -0.68)$

Note that the local maximum at $x \approx 2.88$ is also an absolute maximum.

(f) $(0, 1)$ and $(2, 9)$


9. $y' = \frac{2}{5}x^{-4/5}$

| | | |
|-----------------|------------|------------|
| Intervals | $x < 0$ | $0 < x$ |
| Sign of y' | + | + |
| Behavior of y | Increasing | Increasing |

$y'' = -\frac{8}{25}x^{-9/5}$

| | | |
|-----------------|------------|--------------|
| Intervals | $x < 0$ | $0 < x$ |
| Sign of y'' | + | - |
| Behavior of y | Concave up | Concave down |

Graphical support:



$[-6, 6]$ by $[-1.5, 7.5]$

- (a) $(-\infty, \infty)$
- (b) None
- (c) $(-\infty, 0)$
- (d) $(0, \infty)$
- (e) None
- (f) $(0, 3)$


10. $y' = -\frac{1}{3}x^{-2/3}$

| | | |
|-----------------|------------|------------|
| Intervals | $x < 0$ | $0 < x$ |
| Sign of y' | - | - |
| Behavior of y | Decreasing | Decreasing |

$y'' = \frac{2}{9}x^{-5/3}$

| | | |
|-----------------|--------------|------------|
| Intervals | $x < 0$ | $0 < x$ |
| Sign of y'' | - | + |
| Behavior of y | Concave down | Concave up |

Graphical support:



$[-8, 8]$ by $[0, 10]$

- (a) $(0, \infty)$
- (b) $(-\infty, 0)$


11. $y' = \begin{cases} 2, & x < 1 \\ -2x, & x > 1 \end{cases}$

| | | |
|-----------------|------------|------------|
| Intervals | $x < 1$ | $1 < x$ |
| Sign of y' | + | - |
| Behavior of y | Increasing | Decreasing |

$y'' = \begin{cases} 0, & x < 1 \\ -2, & x > 1 \end{cases}$

| | | |
|-----------------|---------|--------------|
| Intervals | $x < 1$ | $1 < x$ |
| Sign of y'' | 0 | - |
| Behavior of y | Linear | Concave down |

Graphical support:




$[-2, 3]$ by $[-5, 3]$

- (a) None
- (b) $(1, \infty)$

12. $y' = e^x$
 $y'' = e^x$

Since y' and y'' are both positive on the entire domain, y is increasing and concave up on the entire domain.

Graphical support:



$[0, 2\pi]$ by $[0, 20]$

- (a) $(0, 2\pi)$
- (b) None

13. $y = xe^x$

$$y' = e^x + xe^x$$

| | | |
|-----------------|------------|------------|
| Intervals | $x < -1$ | $x > -1$ |
| Sign of y' | - | + |
| Behavior of y | Decreasing | Increasing |

$$y'' = 2e^x + xe^x$$

| | | |
|-----------------|--------------|------------|
| Intervals | $x < -2$ | $x > -2$ |
| Sign of y'' | - | + |
| Behavior of y | Concave down | Concave up |

$$\left(-2, -\frac{2}{e^2}\right)$$

14. $y = x\sqrt{9-x^2}$

$$y' = \sqrt{9-x^2} - \frac{x^2}{\sqrt{9-x^2}} = 0$$

$$x = \pm \frac{3\sqrt{2}}{2}$$

| | | | |
|-----------------|---------------------------------|--------------------------------------------------|-------------------------------|
| Intervals | $-3 < x < -\frac{3\sqrt{2}}{2}$ | $-\frac{3\sqrt{2}}{2} < x < \frac{3\sqrt{2}}{2}$ | $\frac{3\sqrt{2}}{2} < x < 3$ |
| Sign of y' | - | + | - |
| Behavior of y | Decreasing | Increasing | Decreasing |

$$y'' = -\frac{3x}{(9-x^2)^{3/2}} + \frac{x^3}{(9-x^2)^{5/2}} = 0$$

$$y'' = 0 \text{ at } x = 0$$

| | | |
|-----------------|--------------|--------------|
| Intervals | $-3 < x < 0$ | $0 < x < 3$ |
| Sign of y'' | + | - |
| Behavior of y | Concave up | Concave down |

15. $y' = \frac{1}{1+x^2}$

since $y' > 0$ for all x , y is always increasing:

$$y'' = \frac{d}{dx}(1+x^2)^{-1} = -(1+x^2)^{-2}(2x) = \frac{-2x}{(1+x^2)^2}$$

| | | |
|-----------------|------------|--------------|
| Intervals | $x < 0$ | $0 < x$ |
| Sign of y'' | + | - |
| Behavior of y | Concave up | Concave down |

(0, 0)

16. $y = x^3(4-x)$

$$y' = 12x^2 - 4x^3$$

| | | | |
|-----------------|------------|-------------|------------|
| Intervals | $x < 0$ | $0 < x < 3$ | $x > 3$ |
| Sign of y' | + | + | - |
| Behavior of y | Increasing | Increasing | Decreasing |

$$y'' = 24x - 12x^2$$

| | | | |
|-----------------|--------------|-------------|--------------|
| Intervals | $x < 0$ | $0 < x < 2$ | $x > 2$ |
| Sign of y'' | - | + | - |
| Behavior of y | Concave down | Concave up | Concave down |

(0, 0) and (2, 16)

17. $y = x^{1/3}(x-4) = x^{4/3} - 4x^{1/3}$

$$y' = \frac{4}{3}x^{1/3} - \frac{4}{3}x^{-2/3} = \frac{4x-4}{3x^{2/3}}$$

| | | | |
|-----------------|------------|-------------|------------|
| Intervals | $x < 0$ | $0 < x < 1$ | $1 < x$ |
| Sign of y' | - | - | + |
| Behavior of y | Decreasing | Decreasing | Increasing |

$$y'' = \frac{4}{9}x^{-2/3} + \frac{8}{9}x^{-5/3} = \frac{4x+8}{9x^{5/3}}$$

| | | | |
|-----------------|------------|--------------|------------|
| Intervals | $x < -2$ | $-2 < x < 0$ | $0 < x$ |
| Sign of y'' | + | - | + |
| Behavior of y | Concave up | Concave down | Concave up |

 $(-2, 6\sqrt[3]{2}) \approx (-2, 7.56)$ and (0, 0)


18. $y = x^{1/2}(x+3)$

 $y' = \frac{1}{2}x^{-1/2}(x+3) + x^{1/2}$ y is always increasing, so there are no critical points for y' .

$$y'' = \frac{1}{(x)^{3/2}} - \frac{x-3}{4(x)^{5/2}} = 0$$

| | | |
|-----------------|-------------|--------------|
| Intervals | $0 < x < 1$ | $x > 1$ |
| Sign of y'' | + | - |
| Behavior of y | Concave up | Concave down |

(1, 4)

19. We use a combination of analytic and grapher techniques to solve this problem. Depending on the viewing window chosen, graphs obtained using NDER may exhibit strange behavior near $x = 2$ because, for example, NDER $(y, 2) \approx 1,000,000$ while y' is actually undefined at $x = 2$. The graph of $y = \frac{x^3 - 2x^2 + x - 1}{x - 2}$ is shown below.


[-4.7, 4.7] by [-5, 15]

19. Continued

$$y' = \frac{(x-2)(3x^2 - 4x + 1) - (x^3 - 2x^2 + x - 1)(1)}{(x-2)^2}$$

$$= \frac{2x^3 - 8x^2 + 8x - 1}{(x-2)^2}$$

The graph of y' is shown below.



$[-4.7, 4.7]$ by $[-10, 10]$

The zeros of y' are $x \approx 0.15$, $x = 1.40$, and $x = 2.45$.

| Intervals | $x < 0.15$ | $0.15 < x < 1.40$ | $1.40 < x < 2$ | $2 < x < 2.45$ | $2.45 < x$ |
|-----------------|------------|-------------------|----------------|----------------|------------|
| Sign of y' | - | + | - | - | + |
| Behavior of y | Decreasing | Increasing | Decreasing | Decreasing | Increasing |


$$y'' = \frac{(x-2)^2(6x^2 - 16x + 8) - (2x^3 - 8x^2 + 8x - 1)(2)(x-2)}{(x-2)^4}$$

$$= \frac{(x-2)(6x^2 - 16x + 8) - 2(2x^3 - 8x^2 + 8x - 1)}{(x-2)^3}$$

$$= \frac{2x^3 - 12x^2 + 24x - 14}{(x-2)^3}$$

$$= \frac{2(x-1)(x^2 - 5x + 7)}{(x-2)^3}$$

The graph of y'' is shown below.



$[-4.7, 4.7]$ by $[-10, 10]$

Note that the discriminant of $x^2 - 5x + 7$ is

$$(-5)^2 - 4(1)(7) = -3, \text{ so the only solution of } y'' = 0 \text{ is } x = 1.$$

| Intervals | $x < 1$ | $1 < x < 2$ | $2 < x$ |
|-----------------|------------|--------------|------------|
| Sign of y'' | + | - | + |
| Behavior of y | Concave up | Concave down | Concave up |

$(1, 1)$

20. $y' = \frac{(x^2+1)(1) - x(2x)}{(x^2+1)^2} = \frac{-x^2+1}{(x^2+1)^2}$

| Intervals | $x < -1$ | $-1 < x < 1$ | $1 < x$ |
|-----------------|------------|--------------|------------|
| Sign of y' | - | + | - |
| Behavior of y | Decreasing | Increasing | Decreasing |

$$y'' = \frac{(x^2+1)^2(-2x) - (-x^2+1)(2)(x^2+1)(2x)}{(x^2+1)^4}$$

$$= \frac{(x^2+1)(-2x) - 4x(-x^2+1)}{(x^2+1)^3}$$

$$= \frac{2x^3 - 6x}{(x^2+1)^3} = \frac{2x(x^2 - 3)}{(x^2+1)^3}$$

| Intervals | $x < -\sqrt{3}$ | $-\sqrt{3} < x < 0$ | $0 < x < \sqrt{3}$ | $\sqrt{3} < x$ |
|-----------------|-----------------|---------------------|--------------------|----------------|
| Sign of y'' | - | + | - | + |
| Behavior of y | Concave down | Concave up | Concave down | Concave up |

$(0, 0)$, $(\sqrt{3}, \frac{\sqrt{3}}{4})$, and $(-\sqrt{3}, -\frac{\sqrt{3}}{4})$

21. (a) Zero: $x = \pm 1$;
 positive: $(-\infty, -1)$ and $(1, \infty)$;
 negative: $(-1, 1)$
 (b) Zero: $x = 0$;
 positive: $(0, \infty)$;
 negative: $(-\infty, 0)$
22. (a) Zero: $x \approx 0, \pm 1.25$;
 positive: $(-1.25, 0)$ and $(1.25, \infty)$;
 negative: $(-\infty, -1.25)$ and $(0, 1.25)$
 (b) Zero: $x \approx \pm 0.7$;
 positive: $(-\infty, -0.7)$ and $(0.7, \infty)$;
 negative: $(-0.7, 0.7)$
23. (a) $(-\infty, -2]$ and $[0, 2]$
 (b) $[-2, 0]$ and $[2, \infty)$
 (c) Local maxima: $x = -2$ and $x = 2$;
 local minimum: $x = 0$
24. (a) $[-2, 2]$
 (b) $(-\infty, -2]$ and $[2, \infty)$
 (c) Local maximum: $x = 2$;
 local minimum: $x = -2$
25. (a) $v(t) = x'(t) = 2t - 4$
 (b) $a(t) = v'(t) = 2$
 (c) It begins at position 3 moving in a negative direction. It moves to position -1 when $t = 2$, and then changes direction, moving in a positive direction thereafter.
26. (a) $v(t) = x'(t) = -2 - 2t$
 (b) $a(t) = v'(t) = -2$
 (c) It begins at position 6 and moves in the negative direction thereafter.
27. (a) $v(t) = x'(t) = 3t^2 - 3$
 (b) $a(t) = v'(t) = 6t$

27. Continued

- (c) It begins at position 3 moving in a negative direction. It moves to position 1 when $t = 1$, and then changes direction, moving in a positive direction thereafter.

28. (a) $v(t) = x'(t) = 6t - 6t^2$

(b) $a(t) = v'(t) = 6 - 12t$

- (c) It begins at position 0. It starts moving in the positive direction until it reaches position 1 when $t = 1$, and then it changes direction. It moves in the negative direction thereafter.

29. (a) The velocity is zero when the tangent line is horizontal, at approximately $t = 2.2$, $t = 6$ and $t = 9.8$.


- (b) The acceleration is zero at the inflection points, approximately $t = 4$, $t = 8$ and $t = 11$.

30. (a) The velocity is zero when the tangent line is horizontal, at approximately $t = -0.2$, $t = 4$, and $t = 12$.

- (b) The acceleration is zero at the inflection points, approximately $t = 1.5$, $t = 5.2$, $t = 8$, $t = 11$, and $t = 13$.

31. Some calculators use different logistic regression equations, so answers may vary.

(a) $y = \frac{12655.179}{1 + 12.871e^{-0.0326t}}$




(c) $y = \frac{12655.179}{1 + 12.871e^{-0.0326(180)}} = 12,209,870$. (This is remarkably close to the 2000 census number of 12,281,054.)

- (d) The second derivative has a zero at about 78, indicating that the population was growing fastest in 1898. This corresponds to the inflection point on the regression curve.

- (e) The regression equation predicts a population limit of about 12,655,179.

32. Some calculators use different logistic regression equations, so answers may vary.

(a) $y = \frac{28984386.288}{1 + 49.252e^{-0.851t}}$



- (c) The zero of the second derivative is about 4.6, which puts the fastest growth during 1981. This corresponds to the inflection point on the regression curve.

- (d) The regression curve predicts that cable subscribers will approach a limit of $28,984,386 + 12,168,450$ subscribers (about 41 million).

33. $y = 3x - x^3 + 5$

$$y' = 3 - 3x^2$$

$$y'' = -6x$$

$$y' = 0 \text{ at } \pm 1.$$

- $y''(-1) > 0$ and $y''(1) < 0$, so there is a local minimum at $(-1, 3)$ and a local maximum at $(1, 7)$.

34. $y = x^5 - 80x + 100$

$$y' = 5x^4 - 80$$

$$y'' = 20x^3$$

$$y' = 0 \text{ at } \pm 2$$

- $y''(-2) < 0$ and $y''(2) > 0$, so there is a local maximum at $(-2, 228)$ and a local minimum at $(2, -28)$.

35. $y = x^3 + 3x^2 - 2$

$$y' = 3x^2 + 6x$$

$$y'' = 6x + 6$$

$$y' = 0 \text{ at } -2 \text{ and } 0.$$

$$y''(-2) < 0, y''(0) > 0,$$

- so there is a local maximum at $(-2, 2)$ and a local minimum at $(0, -2)$.

36. $y = 3x^5 - 25x^3 + 60x + 20$

$$y' = 15x^4 - 75x^2 + 60$$

$$y'' = 60x^3 - 150x$$

$$y' = 0 \text{ at } \pm 1 \text{ and } \pm 2.$$

$$y''(-2) < 0, y''(-1) > 0$$

$$y''(1) < 0, \text{ and } y''(2) > 0;$$

- so there are local maxima at $(-2, 4)$ and $(1, 58)$, and there are local minima at $(-1, -18)$ and $(2, 36)$.

37. $y = xe^x$

$$y' = (x + 1)e^x$$

$$y'' = (x + 2)e^x$$

$$y' = 0 \text{ at } -1.$$

- $y''(-1) > 0$, so there is a local minimum at $(-1, -1/e)$.

38. $y = xe^{-x}$

$$y' = (1 - x)e^{-x}$$

$$y'' = (x - 2)e^{-x}$$

$$y' = 0 \text{ at } 1$$

- $y''(1) < 0$, so there is a local maximum at $(1, 1/e)$.

39. $y' = (x - 1)^2(x - 2)$

| Intervals | $x < 1$ | $1 < x < 2$ | $2 < x$ |
|-----------------|------------|-------------|------------|
| Sign of y' | - | - | + |
| Behavior of y | Decreasing | Decreasing | Increasing |

39. Continued

$$\begin{aligned}
 y'' &= (x-1)^2(1) + (x-2)(2)(x-1) \\
 &= (x-1)[(x-1) + 2(x-2)] \\
 &= (x-1)(3x-5)
 \end{aligned}$$

| | | | |
|-----------------|------------|-----------------------|-------------------|
| Intervals | $x < 1$ | $1 < x < \frac{5}{3}$ | $\frac{5}{3} < x$ |
| Sign of y'' | + | - | + |
| Behavior of y | Concave up | Concave down | Concave up |

(a) There are no local maxima.

(b) There is a local (and absolute) minimum at $x = 2$.

(c) There are points of inflection at $x = 1$ and at $x = \frac{5}{3}$.

40. $y' = (x-1)(x-2)(x-4)$


| | | | | |
|-----------------|------------|-------------|-------------|------------|
| Intervals | $x < 1$ | $1 < x < 2$ | $2 < x < 4$ | $4 < x$ |
| Sign of y' | + | + | - | + |
| Behavior of y | Increasing | Increasing | Decreasing | Increasing |

$$\begin{aligned}
 y'' &= \frac{d}{dx} [(x-1)^2(x^2 - 6x + 8)] \\
 &= (x-1)^2(2x-6) + (x^2 - 6x + 8)(2)(x-1) \\
 &= (x-1)[(x-1)(2x-6) + 2(x^2 - 6x + 8)] \\
 &= (x-1)(4x^2 - 20x + 22) \\
 &= 2(x-1)(2x^2 - 10x + 11)
 \end{aligned}$$

Note that the zeros of y'' are $x = 1$ and

$$\begin{aligned}
 x &= \frac{10 \pm \sqrt{10^2 - 4(2)(11)}}{4} = \frac{10 \pm \sqrt{12}}{4} \\
 &= \frac{5 \pm \sqrt{3}}{2} \approx 1.63 \text{ or } 3.37.
 \end{aligned}$$

The zeros of y'' can also be found graphically, as shown.





$[-3, 7]$ by $[-8, 4]$

| | | | | |
|-----------------|--------------|----------------|-------------------|------------|
| Intervals | $x < 1$ | $1 < x < 1.63$ | $1.63 < x < 3.37$ | $3.37 < x$ |
| Sign of y'' | - | + | - | + |
| Behavior of y | Concave down | Concave up | Concave down | Concave up |

(a) Local maximum at $x = 2$

(b) Local minimum at $x = 4$


(c) Points of inflection at $x = 1$, at $x \approx 1.63$, and at $x \approx 3.37$.




43. No f must have a horizontal tangent at that point, but f could be increasing (or decreasing), and there would be no local extremum. For example, if $f(x) = x^3$, $f'(0) = 0$ but there is no local extremum at $x = 0$.

44. No. $f''(x)$ could still be positive (or negative) on both sides of $x = c$, in which case the concavity of the function would not change at $x = c$. For example, if $f(x) = x^4$, then $f''(0) = 0$, but f has no inflection point at $x = 0$.


45. One possible answer:




46. One possible answer:



47. One possible answer:



48. One possible answer:



49. (a) $[0, 1]$, $[3, 4]$, and $[5.5, 6]$

(b) $[1, 3]$ and $[4, 5.5]$

(c) Local maxima: $x = 1, x = 4$
 (if f is continuous at $x = 4$), and $x = 6$;
 local minima: $x = 0, x = 3$, and $x = 5.5$

50. If f is continuous on the interval $[0, 3]$:

(a) $[0, 3]$


(b) Nowhere

(c) Local maximum: $x = 3$;
 local minimum: $x = 0$

51. (a) Absolute maximum at $(1, 2)$;
 absolute minimum at $(3, -2)$

(b) None


(c) One possible answer:





52. (a) Absolute maximum at $(0, 2)$;
 absolute minimum at $(2, -1)$ and $(-2, -1)$

(b) At $(1, 0)$ and $(-1, 0)$

(c) One possible answer:



(d) Since f is even, we know $f(3) = f(-3)$. By the continuity of f , since $f(x) < 0$ when $2 < x < 3$, we know that $f(3) \leq 0$, and since $f(2) = -1$ and $f'(x) > 0$ when $2 < x < 3$, we know that $f(3) > -1$. In summary, we know that $f(3) = f(-3), -1 < f(3) \leq 0$, and $-1 < f(-3) \leq 0$.



55. False. For example, consider $f(x) = x^4$ at $c = 0$.

56. True. This is the Second Derivative Test for a local maximum.

57. A. $y = ax^3 + 3x^2 = 4x + 5$ say $a = -2$

$$y' = -6x^2 + 6x + 4$$

$$y'' = -12x + 6$$

$$y'' = 0 \text{ at } \frac{1}{2}$$

| Interval | $x < 1/2$ | $x > 1/2$ |
|-----------------|------------|--------------|
| Sign of y'' | + | - |
| Behavior of y | Concave up | Concave down |

58. E.

59. C. $y = x^5 - 5x^4 + 3x + 7$

$$y' = 5x^4 - 20x^3 + 3$$

$$y'' = 20x^3 - 60x^2$$

$$y'' = 0 \text{ at } 3$$

| Interval | $x < 3$ | $x > 3$ |
|-----------------|--------------|------------|
| Sign of y'' | - | + |
| Behavior of y | Concave down | Concave up |

3 is an inflection point.

60. A.

61. (a) In exercise 13, $a = 4$ and $b = 21$, so $-\frac{b}{3a} = -\frac{7}{4}$, which is the x -value where the point of inflection occurs. The local extrema are at $x = -2$ and $x = -\frac{3}{2}$, which are symmetric about $x = -\frac{7}{4}$.

(b) In exercise 8, $a = -2$ and $b = 6$, so $-\frac{b}{3a} = 1$, which is the x -value where the point of inflection occurs. The local extrema are at $x = 0$ and $x = 2$, which are symmetric about $x = 1$.

(c) $f'(x) = 3ax^2 + 2bx + c$ and
 $f''(x) = 6ax + 2b$.

The point of inflection will occur where

$$f''(x) = 0, \text{ which is at } x = -\frac{b}{3a}.$$

If there are local extrema, they will occur at the zeros of $f'(x)$. Since $f'(x)$ is quadratic, its graph is a parabola and any zeros will be symmetric about the vertex which will also be where $f''(x) = 0$.

$$\begin{aligned} 62. (a) f'(x) &= \frac{(1 + ae^{-bx})(0) - (c)(-abe^{-bx})}{(1 + ae^{-bx})^2} \\ &= \frac{abce^{-bx}}{(1 + ae^{-bx})^2} \\ &= \frac{abce^{-bx}}{(e^{bx} + a)^2}, \end{aligned}$$

so the sign of $f'(x)$ is the same as the sign of abc .

$$\begin{aligned} (b) f''(x) &= \frac{(e^{bx} + a)^2(ab^2ce^{bx}) - (abce^{bx})2(e^{bx} + a)(be^{bx})}{(e^{bx} + a)^4} \\ &= \frac{(e^{bx} + a)(ab^2ce^{bx}) - (abce^{bx})(2be^{bx})}{(e^{bx} + a)^3} \\ &= -\frac{ab^2ce^{bx}(e^{bx} - a)}{(e^{bx} + a)^3} \end{aligned}$$

Since $a > 0$, this changes sign when $x = \frac{\ln a}{b}$ due to the $e^{bx} - a$ factor in the numerator, and $f(x)$ has a point of inflection at the location.

63. (a) $f'(x) = 4ax^3 + 3bx^2 + 2cx + d$

$$f''(x) = 12ax^2 + 6bx + 2c$$

Since $f''(x)$ is quadratic, it must have 0, 1, or 2 zeros. If $f''(x)$ has 0 or 1 zeros, it will not change sign and the concavity of $f(x)$ will not change, so there is no point of inflection. If $f''(x)$ has 2 zeros, it will change sign twice, and $f(x)$ will have 2 points of inflection.

(b) If f has no points of inflection, then $f''(x)$ has 0 or 1 zeros, so the discriminant of $f''(x)$ is ≤ 0 . This gives $(6b)^2 - 4(12a)(2c) \leq 0$, or $3b^2 \leq 8ac$. If f has 2 points of inflection, then $f''(x)$ has 2 zeros and the inequality is reversed, so $3b^2 > 8ac$. In summary, f has 2 points of inflection if and only if $3b^2 > 8ac$.

Quick Quiz Sections 4.1-4.3

1. (C) $f'(x) = 5(x-2)^4(x+3)^4 + 4(x-2)^5(x+3)^3 = 0$

$$x = -3, -\frac{7}{9}, 2$$

2. (D) $f'(x) = (x-3)^2 + 2(x-2)(x-3) = 0$

$$f'(x) = (x-3)(3x-7) = 0$$

$$x = \frac{7}{3}, 3$$

3. (B) $x^2 - 9 = 0$

$$x = \pm 3$$

4. (a) $\frac{d}{dx} 3 \ln(x^2 + 2) - 2x$

$$= 3 \frac{2x}{x^2 + 2} - 2 = 0$$

$$x = 1, 2$$

| Intervals | $-2 < x < 1$ | $1 < x < 2$ | $2 < x < 4$ |
|-----------------|--------------|-------------|-------------|
| Sign of y' | - | + | - |
| Behavior of y | Decreasing | Increasing | Decreasing |

f has relative minima at $x = 1$ and $x = 4$ f has relative maxima at $x = \pm 2$

(b) $f''(x) = \frac{d}{dx} \left(\frac{6x}{x^2 + 2} - 2 \right)$

$$f''(x) = \frac{6}{x^2 + 2} - \frac{12x^2}{(x^2 + 2)^2} = 0$$

$$x = \pm\sqrt{2}$$

 f has points of inflection at $x = \pm\sqrt{2}$

(c) The absolute maximum is at $x = -2$ and $f(x) = 3 \ln 6 + 4$.

Section 4.4 Modeling and Optimization (pp. 219–232)

Exploration 1 Constructing Cones


1. The circumference of the base of the cone is the circumference of the circle of radius 4 minus x , or $8\pi - x$.

Thus, $r = \frac{8\pi - x}{2\pi}$. Use the Pythagorean Theorem to find h , and the formula for the volume of a cone to find V .

2. The expression under the radical must be nonnegative, that


$$\text{is, } 16 - \left(\frac{8\pi - x}{2\pi}\right)^2 \geq 0.$$

Solving this inequality for x gives: $0 \leq x \leq 16\pi$.



$[0, 16\pi]$ by $[-10, 40]$

3. The circumference of the original circle of radius 4 is 8π . Thus, $0 \leq x \leq 8\pi$.



$[0, 8\pi]$ by $[-10, 40]$

4. The maximum occurs at about $x = 4.61$. The maximum volume is about $V = 25.80$.

5. Start with $\frac{dV}{dx} = \frac{2\pi}{3}rh \frac{dr}{dx} + \frac{\pi}{3}r^2 \frac{dh}{dx}$.

Compute $\frac{dr}{dx}$ and $\frac{dh}{dx}$, substitute these values in

$\frac{dV}{dx}$, set $\frac{dV}{dx} = 0$, and solve for x to obtain

$$x = \frac{8(3 - \sqrt{6})\pi}{3} \approx 4.61.$$

$$\text{Then } V = \frac{128\pi\sqrt{3}}{27} \approx 25.80.$$

Quick Review 4.4

1. $y' = 3x^2 - 12x + 12 = 3(x - 2)^2$

Since $y' \geq 0$ for all x (and $y' > 0$ for $x \neq 2$), y is increasing on $(-\infty, \infty)$ and there are no local extrema.

2. $y' = 6x^2 + 6x - 12 = 6(x + 2)(x - 1)$

$$y'' = 12x + 6$$

The critical points occur at $x = -2$ or $x = 1$, since $y' = 0$ at these points. Since $y''(-2) = -18 < 0$, the graph has a local maximum at $x = -2$. Since $y''(1) = 18 > 0$, the graph has a

local minimum at $x = 1$. In summary, there is a local maximum at $(-2, 17)$ and a local minimum at $(1, -10)$.

3. $V = \frac{1}{3}\pi r^2 h = \frac{1}{3}\pi(5)^2(8) = \frac{200\pi}{3} \text{ cm}^3$

4. $V = \pi r^2 h = 1000$

$$SA = 2\pi r h + 2\pi r^2 = 600$$

Solving the volume equation for h gives $h = \frac{1000}{\pi r^2}$.

Substituting into the surface area equation gives $\frac{2000}{r} + 2\pi r^2 = 600$. Solving graphically, we have

$r \approx -11.14$, $r \approx 4.01$, or $r \approx 7.13$. Discarding the negative

value and using $h = \frac{1000}{\pi r^2}$ to find the corresponding values

of h , the two possibilities for the dimensions of the cylinder are:

$r \approx 4.01$ cm and $h \approx 19.82$ cm, or,

$r \approx 7.13$ cm and $h \approx 6.26$ cm.

5. Since $y = \sin x$ is an odd function, $\sin(-\alpha) = -\sin \alpha$.

6. Since $y = \cos x$ is an even function, $\cos(-\alpha) = \cos \alpha$.

7. $\sin(\pi - \alpha) = \sin \pi \cos \alpha - \cos \pi \sin \alpha$
 $= 0 \cos \alpha - (-1) \sin \alpha$
 $= \sin \alpha$

8. $\cos(\pi - \alpha) = \cos \pi \cos \alpha - \sin \pi \sin \alpha$
 $= (-1) \cos \alpha + 0 \sin \alpha$
 $= -\cos \alpha$

9. $x^2 + y^2 = 4$ and $y = \sqrt{3}x$

$$x^2 + (\sqrt{3}x)^2 = 4$$

$$x^2 + 3x^2 = 4$$

$$4x^2 = 4$$

$$x = \pm 1$$

Since $y = \sqrt{3}x$, the solution are:

$$x = 1 \text{ and } y = \sqrt{3}, \text{ or } x = -1 \text{ and } y = -\sqrt{3}.$$

In ordered pair notation, the solutions are

$$(1, \sqrt{3}) \text{ and } (-1, -\sqrt{3}).$$

10. $\frac{x^2}{4} + \frac{y^2}{9} = 1$ and $y = x + 3$

$$\frac{x^2}{4} + \frac{(x+3)^2}{9} = 1$$

$$9x^2 + 4(x+3)^2 = 36$$

$$9x^2 + 4x^2 + 24x + 36 = 36$$

$$13x^2 + 24x = 0$$

$$x(13x + 24) = 0$$

$$x = 0 \text{ or } x = -\frac{24}{13}$$

10. Continued

Since $y = x + 3$, the solutions are:

$$x = 0 \text{ and } y = 3, \text{ or } x = -\frac{24}{13} \text{ and } y = \frac{15}{13}.$$

In ordered pair notation, the solutions are $(0, 3)$ and

$$\left(-\frac{24}{13}, \frac{15}{13}\right).$$

Section 4.4 Exercises


1. Represent the numbers by x and $20 - x$, where $0 \leq x \leq 20$.

(a) The sum of the squares is given by

$$f(x) = x^2 + (20 - x)^2 = 2x^2 - 40x + 400. \text{ Then}$$

$f'(x) = 4x - 40$. The critical point and endpoints occur at $x = 0$, $x = 10$, and $x = 20$. Then $f(0) = 400$, $f(10) = 200$, and $f(20) = 400$. The sum of the squares is as large as possible for the numbers 0 and 20, and is as small as possible for the numbers 10 and 10.

Graphical support:



$[0, 20]$ by $[0, 450]$

(b) The sum of one number plus the square root of the other is given by $g(x) = x + \sqrt{20 - x}$. Then

$$g'(x) = 1 - \frac{1}{2\sqrt{20 - x}}. \text{ The critical point occurs when}$$

$$2\sqrt{20 - x} = 1, \text{ so } 20 - x = \frac{1}{4} \text{ and } x = \frac{79}{4}.$$

Testing the endpoints and critical point, we find $g(0) = \sqrt{20} \approx$

$$4.47, g\left(\frac{79}{4}\right) = \frac{81}{4} = 20.25, \text{ and } g(20) = 20. \text{ The sum is}$$


as large as possible when the numbers are

$$\frac{79}{4} \text{ and } \frac{1}{4} \left(\text{summing } \frac{79}{4} + \sqrt{\frac{1}{4}}\right), \text{ and is as small as}$$


possible when the numbers are 0 and 20

$$\left(\text{summing } 0 + \sqrt{20}\right).$$

Graphical support:



$[0, 20]$ by $[-10, 25]$



3. Let x represent the length of the rectangle in inches ($x > 0$).

Then the width is $\frac{16}{x}$ and the perimeter is

$$P(x) = 2\left(x + \frac{16}{x}\right) = 2x + \frac{32}{x}.$$

Since $P'(x) = 2 - 32x^{-2} = \frac{2(x^2 - 16)}{x^2}$ this critical point

occurs at $x = 4$. Since $P'(x) < 0$ for $0 < x < 4$ and $P'(x) > 0$ for $x > 4$, this critical point corresponds to the minimum perimeter. The smallest possible perimeter is $P(4) = 16$ in., and the rectangle's dimensions are 4 in. by 4 in.

Graphical support:



$[0, 20]$ by $[0, 40]$



5. (a) The equation of line AB is $y = -x + 1$, so the y -coordinate of P is $-x + 1$.

(b) $A(x) = 2x(1 - x)$

(c) Since $A'(x) = \frac{d}{dx}(2x - 2x^2) = 2 - 4x$, the critical point occurs at $x = \frac{1}{2}$. Since $A'(x) > 0$ for $0 < x < \frac{1}{2}$ and


$A'(x) < 0$ for $\frac{1}{2} < x < 1$, this critical point corresponds

to the maximum area. The largest possible area is


$A\left(\frac{1}{2}\right) = \frac{1}{2}$ square unit, and the dimensions of the

rectangle are $\frac{1}{2}$ unit by 1 unit.

Graphical support:



$[0, 1]$ by $[-0.5, 1]$



7. Let x be the side length of the cut-out square ($0 < x < 4$). Then the base measures $8 - 2x$ in. by $15 - 2x$ in., and the volume is

$V(x) = x(8 - 2x)(15 - 2x) = 4x^3 - 46x^2 + 120x$. Then

$V'(x) = 12x^2 - 92x + 120 = 4(3x - 5)(x - 6)$.

Then the critical point (in $0 < x < 4$) occurs at $x = \frac{5}{3}$. Since


$V'(x) > 0$ for $0 < x < \frac{5}{3}$ and $V'(x) < 0$ for $\frac{5}{3} < x < 4$,

the critical point corresponds to the maximum volume.


The maximum volume is $V\left(\frac{5}{3}\right) = \frac{2450}{27} \approx 90.74 \text{ in}^3$, and the


dimensions are $\frac{5}{3}$ in. by $\frac{14}{3}$ in. by $\frac{35}{3}$ in.

Graphical support:



$[0, 4]$ by $[-25, 100]$





9. Let x be the length in meters of each side that adjoins the river. Then the side parallel to the river measures $800 - 2x$ meters and the area is


$$A(x) = x(800 - 2x) = 800x - 2x^2 \text{ for } 0 < x < 400.$$

Therefore, $A'(x) = 800 - 4x$ and the critical point occurs at $x = 200$. Since $A'(x) > 0$ for $0 < x < 200$ and

$A'(x) < 0$ for $200 < x < 400$, the critical point corresponds to the maximum area. The largest possible area is

$A(200) = 80,000 \text{ m}^2$ and the dimensions are 200 m (perpendicular to the river) by 400 m (parallel to the river).

Graphical support:



$[0, 400]$ by $[-25,000, 90,000]$




11. (a) Let x be the length in feet of each side of the square base. Then the height is $\frac{500}{x^2}$ ft and the surface area (not including the open top) is

$$S(x) = x^2 + 4x\left(\frac{500}{x^2}\right) = x^2 + 2000x^{-1}. \text{ Therefore,}$$


$$S'(x) = 2x - 2000x^{-2} = \frac{2(x^3 - 1000)}{x^2} \text{ and the critical}$$

point occurs at $x = 10$. Since $S'(x) < 0$ for $0 < x < 10$ and $S'(x) > 0$ for $x > 10$, the critical point corresponds to the minimum amount of steel used. The dimensions should be 10 ft by 10 ft by 5 ft, where the height is 5 ft.

(b) Assume that the weight is minimized when the total area of the bottom and the four sides is minimized.



13. Let x be the height in inches of the printed area. Then the width of the printed area is $\frac{50}{x}$ in. and the overall dimensions are $x + 8$ in. by $\frac{50}{x} + 4$ in. The amount of paper used is $A(x) = (x + 8)\left(\frac{50}{x} + 4\right) = 4x + 82 + \frac{400}{x} \text{ in}^2$. Then $A'(x) = 4 - 400x^{-2} = \frac{4(x^2 - 100)}{x^2}$ and the critical point (for $x > 0$) occurs at $x = 10$. Since $A'(x) < 0$ for $0 < x < 10$ and $A'(x) > 0$ for $x > 10$, the critical point corresponds to the minimum amount of paper. Using $x + 8$ and $\frac{50}{x} + 4$ for $x = 10$, the overall dimensions are 18 in. high by 9 in. wide.



15. We assume that a and b are held constant. Then

$$A(\theta) = \frac{1}{2}ab \sin \theta \text{ and } A'(\theta) = \frac{1}{2}ab \cos \theta. \text{ The critical point}$$

(for $0 < \theta < \pi$) occurs at $\theta = \frac{\pi}{2}$. Since $A'(\theta) > 0$

for $0 < \theta < \frac{\pi}{2}$ and $A'(\theta) < 0$ for $\frac{\pi}{2} < \theta < \pi$,

the critical point corresponds to the maximum area. The

angle that maximizes the triangle's area is $\theta = \frac{\pi}{2}$ (or 90°).

17. Note that $\pi r^2 h = 1000$, so $h = \frac{1000}{\pi r^2}$. Then

$$A = 8r^2 + 2\pi r h = 8r^2 + \frac{2000}{r}, \text{ so}$$

$$\frac{dA}{dr} = 16r - 2000r^{-2} = \frac{16(r^3 - 125)}{r^2}. \text{ The critical point}$$

occurs at $r = \sqrt[3]{125} = 5$ cm. Since $\frac{dA}{dr} < 0$ for $0 < r < 5$ and


$\frac{dA}{dr} > 0$ for $r > 5$, the critical point corresponds to the least amount of aluminium used or wasted and hence the most economical can. The dimensions are $r = 5$ cm and $h = \frac{40}{\pi}$,

so the ratio of h to r is $\frac{8}{\pi}$ to 1.

19. (a) The "sides" of the suitcase will measure $24 - 2x$ in. by $18 - 2x$ in. and will be $2x$ in. apart, so the volume formula is


$$V(x) = 2x(24 - 2x)(18 - 2x) = 8x^3 - 168x^2 + 864x.$$

(b) We require $x > 0$, $2x < 18$, and $2x < 24$. Combining these requirements, the domain is the interval $(0, 9)$.



$[0, 9]$ by $[-400, 1600]$

19. Continued



$[0, 9]$ by $[-400, 1600]$

The maximum volume is approximately 1309.95 when $x \approx 3.39$ in.

(d) $V'(x) = 24x^2 - 336x + 864 = 24(x^2 - 14x + 36)$

The critical point is at

$$x = \frac{4 \pm \sqrt{(-14)^2 - 4(1)(36)}}{2(1)} = \frac{14 \pm \sqrt{52}}{2} = 7 \pm \sqrt{13}, \text{ that}$$

is, $x \approx 3.39$ or $x \approx 10.61$. We discard the larger value because it is not in the domain. Since $V''(x) = 24(2x - 14)$, which is negative when $x \approx 3.39$, the critical point corresponds to the maximum volume. The maximum value occurs at $x = 7 - \sqrt{13} \approx 3.39$, which confirms the results in (c).

(e) $8x^3 - 168x^2 + 864x = 1120$
 $8(x^3 - 21x^2 + 108x - 140) = 0$
 $8(x - 2)(x - 5)(x - 14) = 0$

Since 14 is not in the domain, the possible values of x are $x = 2$ in. or $x = 5$ in.

(f) The dimensions of the resulting box are $2x$ in., $(24 - 2x)$ in., and $(18 - 2x)$ in. Each of these measurements must be positive, so that gives the domain of $(0, 9)$

21. If the upper right vertex of the rectangle is located at $(x, 4 \cos 0.5x)$ for $0 < x < \pi$, then the rectangle has width $2x$ and height $4 \cos 0.5x$, so the area is $A(x) = 8x \cos 0.5x$. Then $A'(x) = 8x(-0.5 \sin 0.5x) + 8(\cos 0.5x)(1)$

$$= -4x \sin 0.5x + 8 \cos 0.5x.$$

Solving $A'(x)$ graphically for $0 < x < \pi$, we find that $x \approx 1.72$. Evaluating $2x$ and $4 \cos 0.5x$ for $x \approx 1.72$, the dimensions of the rectangle are approximately 3.44 (width) by 2.61 (height), and the maximum area is approximately 8.98.

23. Set $r'(x) = c'(x): 4x^{-1/2} = 4x$. The only positive critical value is $x = 1$, so profit is maximized at a production level of 1000 units. Note that $(r - c)''(x) = -2(x)^{-3/2} - 4 < 0$ for all positive x , so the Second Derivative Test confirms the Maximum.

25. Set $c'(x) = \frac{c(x)}{x}$: $3x^2 - 20x + 30 = x^2 - 10x + 30$. The only positive solution is $x = 5$, so average cost is minimized at a production level of 5000 units. Note that

$$\frac{d^2}{dx^2} \left(\frac{c(x)}{x} \right) = 2 > 0 \text{ for all positive } x, \text{ so the Second}$$

Derivative Test Confirms the minimum.

27. Revenue: $r(x) = [200 - 2(x - 50)]x = -2x^2 + 300x$

$$\text{Cost: } c(x) = 6000 + 32x$$

$$\text{Profit: } p(x) = r(x) - c(x)$$

$$= -2x^2 + 268x - 6000, 50 \leq x \leq 80$$

Since $p'(x) = -4x + 268 = -4(x - 67)$, the critical point occurs at $x = 67$. This value represents the maximum because $p''(x) = -4$, which is negative for all x in the domain. The maximum profit occurs if 67 people go on the tour.

29. (a) $f'(x)$ is a quadratic polynomial, and as such it can have 0, 1, or 2 zeros. If it has 0 or 1 zeros, then its sign never changes, so $f(x)$ has no local extrema. If $f'(x)$ has 2 zeros, then its sign changes twice, and $f(x)$ has 2 local extrema at those points.

(b) Possible answers:

No local extrema: $y = x^3$;

2 local extrema: $y = x^3 - 3x$

31. Since $2x + 2y = 36$, we know that $y = 18 - x$. In part (a),

the radius is $\frac{x}{2\pi}$ and the height is $18 - x$, and so the volume is given by

$$\pi r^2 h = \pi \left(\frac{x}{2\pi} \right)^2 (18 - x) = \frac{1}{4\pi} x^2 (18 - x).$$

In part (b), the radius is and the height is $18 - x$, and so the volume is given by $\pi r^2 h = \pi x^2 (18 - x)$. Thus, each problem requires us to find the value of x that maximizes $f(x) = x^2(18 - x)$ in the interval $0 < x < 18$, so the two problems have the same answer.

To solve either problem, note that $f(x) = 18x^2 - x^3$ and so $f'(x) = 36x - 3x^2 = -3x(x - 12)$. The critical point occurs at $x = 12$. Since $f'(x) > 0$ for $0 < x < 12$ and $f'(x) < 0$ for $12 < x < 18$, the critical point corresponds to the maximum value of $f(x)$. To maximize the volume in either part (a) or (b), let $x = 12$ cm and $y = 6$ cm.

33. (a) We require $f(x)$ to have a critical point at $x = 2$. Since

$$f'(x) = 2x - ax^{-2}, \text{ we have } f'(2) = 4 - \frac{a}{4} \text{ and so our}$$

requirement is that $4 - \frac{a}{4} = 0$! Therefore, $a = 16$. To

verify that the critical point corresponds to a local minimum, note that we now have $f'(x) = 2x - 16x^{-2}$ and so $f''(x) = 2 + 32x^{-3}$, so $f''(2) = 6$, which is positive as expected. So, use $a = -16$.

- (b) We require $f''(1) = 0$. Since $f'' = 2 + 2ax^{-3}$, we have

$$f''(1) = 2 + 2a, \text{ so our requirement is that } 2 + 2a = 0.$$

Therefore, $a = -1$. To verify that $x = 1$ is in fact an inflection point, note that we now have


$$f''(x) = 2 - 2x^{-3}, \text{ which is negative for } 0 < x < 1 \text{ and positive for } x > 1. \text{ Therefore, the graph of } f \text{ is concave down in the interval } (0, 1) \text{ and concave up in the interval } (1, \infty). \text{ So, use } a = -1.$$

35. (a) Note that $f'(x) = 3x^2 + 2ax + b$. We require $f'(-1) = 0$ and $f'(3) = 0$, which give $3 - 2a + b = 0$ and $27 + 6a + b = 0$. Subtracting the first equation from the second, we have $24 + 8a = 0$ and so $a = -3$. Substituting into the first equation, we have $9 + b = 0$, so $b = -9$. Therefore, our equation for $f(x)$ is $f(x) = x^3 - 3x^2 - 9x$. To verify that we have a local maximum at $x = -1$ and a local minimum at $x = 3$, note that $f'(x) = 3x^2 - 6x - 9 = 3(x + 1)(x - 3)$, which is positive for $x < -1$, negative for $-1 < x < 3$, and positive for $x > 3$. So, use $a = -3$ and $b = -9$.

- (b) Note that $f'(x) = 3x^2 + 2ax + b$ and $f''(x) = 6x + 2a$. We require $f'(4) = 0$ and $f''(1) = 0$, which give $48 + 8a + b = 0$ and $6 + 2a = 0$. By the second equation, $a = -3$, and so the first equation becomes $48 - 24 + b = 0$. Thus $b = -24$. To verify that we have a local minimum at $x = 4$, and an inflection point at $x = 1$, note that we now have $f''(x) = 6x - 6$. Since f'' changes sign at $x = 1$ and is positive at $x = 4$, the desired conditions are satisfied. So, use $a = -3$ and $b = -24$.


Changing the value of k changes the maximum strength, but not the dimensions of the strongest beam. The graphs for different values of k look the same except that the vertical scale is different.

37. (a) Note that $w^2 + d^2 = 12^2$, so $d = \sqrt{144 - w^2}$. Then we may write $S = kwd^2 = kw(144 - w^2) = 144kw - kw^3$ for $0 < w < 12$, so $\frac{dS}{dw} = 144k - 3kw^2 = -3k(w^2 - 48)$. The critical point (for $0 < w < 12$) occurs at $w = \sqrt{48} = 4\sqrt{3}$. Since $\frac{dS}{dw} > 0$ for $0 < w < 4\sqrt{3}$ and $\frac{dS}{dw} < 0$ for $4\sqrt{3} < w < 12$, the critical point corresponds to the maximum strength. The dimensions are $4\sqrt{3}$ in. wide by $4\sqrt{6}$ in. deep.



$[0, 12]$ by $[-100, 800]$

The graph of $S = 144w - w^3$ is shown. The maximum strength shown in the graph occurs at $w = 4\sqrt{3} \approx 6.9$, which agrees with the answer to part (a).



$[0, 12]$ by $[-100, 800]$

The graph of $S = d^2\sqrt{144 - d^2}$ is shown. The maximum strength shown in the graph occurs at $d = 4\sqrt{6} \approx 9.8$, which agrees with the answer to part (a), and its value is the same as the maximum value found in part (b), as expected.

39. (a) $v(t) = s'(t) = -10\pi \sin \pi t$

The speed at time t is $10\pi|\sin \pi t|$. The maximum speed is 10π cm/sec and it occurs at $t = \frac{1}{2}$, $t = \frac{3}{2}$, $t = \frac{5}{2}$, and $t = \frac{7}{2}$ sec. The position at these times is $s = 0$ cm (rest position), and the acceleration $a(t) = v'(t) = -10\pi^2 \cos \pi t$ is 0 cm/sec² at these times.

39. Continued

(b) Since $a(t) = -10\pi^2 \cos \pi t$, the greatest magnitude of the acceleration occurs at $t = 0$, $t = 1$, $t = 2$, $t = 3$, and $t = 4$. At these times, the position of the cart is either $s = -10$ cm or $s = 10$ cm, and the speed of the cart is 0 cm/sec.

41. The square of the distance is

$$D(x) = \left(x - \frac{3}{2}\right)^2 + (\sqrt{x} + 0)^2 = x^2 - 2x + \frac{9}{4},$$

so $D'(x) = 2x - 2$ and the critical point occurs at $x = 1$.

Since $D'(x) < 0$ for $x < 1$ and $D'(x) > 0$ for $x > 1$, the critical point corresponds to the minimum distance. The minimum

distance is $\sqrt{D(1)} = \frac{\sqrt{5}}{2}$.

43. No. Since $f(x)$ is a quadratic function and the coefficient of x^2 is positive, it has an absolute minimum at the point where $f'(x) = 2x - 1 = 0$, and the point is $\left(\frac{1}{2}, \frac{3}{4}\right)$.

45. (a)
$$\begin{aligned} 2 \sin t &= \sin 2t \\ 2 \sin t &= 2 \sin t \cos t \\ 2(\sin t)(1 - \cos t) &= 0 \\ \sin t &= 0 \text{ or } \cos t = 1 \end{aligned}$$

$t = k\pi$, where k is an integer

The masses pass each other whenever t is an integer multiple of π seconds.

(b) The vertical distance between the objects is the absolute value of $f(x) = \sin 2t - 2 \sin t$.

Find the critical points in $[0, 2\pi]$:

$$\begin{aligned} f'(x) &= 2 \cos 2t - 2 \cos t = 0 \\ 2(2 \cos^2 t - 1) - 2 \cos t &= 0 \\ 2(2 \cos^2 t - \cos t - 1) &= 0 \\ 2(2 \cos t + 1)(\cos t - 1) &= 0 \end{aligned}$$

$$\cos t = -\frac{1}{2} \text{ or } \cos t = 1$$

$$t = \frac{2\pi}{3}, \frac{4\pi}{3}, 0, 2\pi$$

The critical points (and endpoints) are $(0, 0)$,

$$\left(\frac{2\pi}{3}, -\frac{3\sqrt{3}}{2}\right), \left(\frac{4\pi}{3}, \frac{3\sqrt{3}}{2}\right), \text{ and } (2\pi, 0)$$

The distance is greatest when $t = \frac{2\pi}{3}$ sec and when

$t = \frac{4\pi}{3}$ sec. The distance at those times is $\frac{3\sqrt{3}}{2}$ meters.

47. The trapezoid has height $(\cos \theta)$ ft and the trapezoid bases measure 1 ft and $(1 + 2 \sin \theta)$ ft, so the volume is given by


$$\begin{aligned} V(\theta) &= \frac{1}{2}(\cos \theta)(1 + 1 + 2 \sin \theta)(20) \\ &= 20(\cos \theta)(1 + \sin \theta). \end{aligned}$$

Find the critical points for $0 \leq \theta < \frac{\pi}{2}$:

$$\begin{aligned} V'(\theta) &= 20(\cos \theta)(\cos \theta) + 20(1 + \sin \theta)(-\sin \theta) = 0 \\ &20 \cos^2 \theta - 20 \sin \theta - 20 \sin^2 \theta = 0 \\ &20(1 - \sin^2 \theta) - 20 \sin \theta - 20 \sin^2 \theta = 0 \\ &\quad -20(2 \sin^2 \theta + \sin \theta - 1) = 0 \\ &\quad -20(2 \sin \theta - 1)(\sin \theta + 1) = 0 \\ &\sin \theta = \frac{1}{2} \text{ or } \sin \theta = -1 \\ &\quad \theta = \frac{\pi}{6} \end{aligned}$$

The critical point is at $\left(\frac{\pi}{6}, 15\sqrt{3}\right)$. Since

$V'(\theta) > 0$ for $0 \leq \theta < \frac{\pi}{6}$ and $V'(\theta) < 0$ for $\frac{\pi}{6} < \theta < \frac{\pi}{2}$, the critical point corresponds to the maximum possible trough volume. The volume is maximized when $\theta = \frac{\pi}{6}$.



49. Since $R = M^2 \left(\frac{C}{2} - \frac{M}{3} \right) = \frac{C}{2}M^2 - \frac{1}{3}M^3$, we have



$$\frac{dR}{dM} = CM - M^2. \text{ Let } f(M) = CM - M^2. \text{ Then}$$

$f'(M) = C - 2M$, and the critical point for f occurs at

$M = \frac{C}{2}$. This value corresponds to a maximum because

$f'(M) > 0$ for $M < \frac{C}{2}$ and $f'(M) < 0$ for $M > \frac{C}{2}$. The value

of M that maximizes $\frac{dR}{dM}$ is $M = \frac{C}{2}$.



51. True. This is guaranteed by the Extreme Value Theorem (Section 4.1).



53. D. $f(x) = x^2(60 - x)$

$$\begin{aligned} f'(x) &= x^2(-1) + (60 - x)(2x) \\ &= -x^2 + 120x - 2x^2 \\ &= -3x^2 + 120x \\ &= -3x(x - 40) \end{aligned}$$

$$\begin{aligned} x = 0 \quad \text{or} \quad x = 40 \\ 60 - x = 60 \quad \quad 60 - x = 20 \end{aligned}$$

$$\begin{aligned} x^2(60 - x) &= 0 \\ (40)^2(20) &= (1600)(20) \\ &= 32,000 \end{aligned}$$



55. B. $A = \frac{1}{2}bh$

$$b^2 + h^2 = 100$$

$$b = \sqrt{100 - h^2}$$


$$A = \frac{h}{2}\sqrt{100 - h^2}$$

$$A' = \frac{\sqrt{100 - h^2}}{2} - \frac{h^2}{2\sqrt{100 - h^2}}$$


$$A' = 0 \text{ when } h = \sqrt{50}$$

$$b = \sqrt{100 - \sqrt{50}^2} = \sqrt{50}$$

$$A_{\max} = \frac{1}{2}\sqrt{50}\sqrt{50} = 25$$



57. Normal



Let P be the foot of the perpendicular from A to the mirror, and Q be the foot of the perpendicular from B to the mirror.

57. Continued

Suppose the light strikes the mirror at point R on the way from A to B . Let:

a = distance from A to P

b = distance from B to Q

c = distance from P to Q

x = distance from P to R

To minimize the time is to minimize the total distance the light travels going from A to B . The total distance is

$$D(x) = \sqrt{x^2 + a^2} + \sqrt{(c-x)^2 + b^2}$$

Then

$$\begin{aligned} D'(x) &= \frac{1}{2\sqrt{x^2 + a^2}}(2x) + \frac{1}{2\sqrt{(c-x)^2 + b^2}}[-2(c-x)] \\ &= \frac{x}{\sqrt{x^2 + a^2}} - \frac{c-x}{\sqrt{(c-x)^2 + b^2}} \end{aligned}$$

Solving $D'(x) = 0$ gives the equation

$$\frac{x}{\sqrt{x^2 + a^2}} = \frac{c-x}{\sqrt{(c-x)^2 + b^2}} \quad \text{which we will refer to as}$$

Equation 1. Squaring both sides, we have:

$$\begin{aligned} \frac{x^2}{x^2 + a^2} &= \frac{(c-x)^2}{(c-x)^2 + b^2} \\ x^2[(c-x)^2 + b^2] &= (c-x)^2(x^2 + a^2) \\ x^2(c-x)^2 + x^2b^2 &= (c-x)^2x^2 + (c-x)^2a^2 \\ x^2b^2 &= (c-x)^2a^2 \\ x^2b^2 &= [c^2 - 2cx + x^2]a^2 \\ 0 &= (a^2 - b^2)x^2 - 2a^2cx + a^2c^2 \\ 0 &= [(a+b)x - ac][(a-b)x - ac] \\ x &= \frac{ac}{a+b} \quad \text{or} \quad x = \frac{ac}{a-b} \end{aligned}$$

Note that the value $x = \frac{ac}{a-b}$ is an extraneous solution because x and $c-x$ have opposite signs for this value. The only critical point occurs at $x = \frac{ac}{a+b}$.

To verify that critical point represents the minimum distance, note that

$$\begin{aligned} D''(x) &= \frac{(\sqrt{x^2 + a^2})(1) - (x)\left(\frac{x}{\sqrt{x^2 + a^2}}\right)}{x^2 + a^2} \\ &\quad - \frac{(\sqrt{(c-x)^2 + b^2})(-1) - (c-x)\left(\frac{-(c-x)}{\sqrt{(c-x)^2 + b^2}}\right)}{(c-x)^2 + b^2} \\ &= \frac{(x^2 + a^2) - x^2}{(x^2 + a^2)^{3/2}} - \frac{-[(c-x)^2 + b^2] + (c-x)^2}{[(c-x)^2 + b^2]^{3/2}} \\ &= \frac{a^2}{(x^2 + a^2)^{3/2}} + \frac{b^2}{[(c-x)^2 + b^2]^{3/2}}, \end{aligned}$$

which is always positive.

We now know that $D(x)$ is minimized when Equation 1 is


true, or, equivalently, $\frac{PR}{AR} = \frac{QR}{BR}$. This means that the two right triangles APR and BQR are similar, which in turn implies that the two angles must be equal.

$$59. (a) \quad v = cr_0r^2 - cr^3$$

$$\frac{dv}{dr} = 2cr_0r - 3cr^2 = cr(2r_0 - 3r)$$

The critical point occurs at $r = \frac{2r_0}{3}$. (Note that $r = 0$ is not in the domain of v .) The critical point represents a maximum because $\frac{d^2v}{dr^2} = 2cr_0 - 6cr = 2c(r_0 - 3r)$, which is negative in the domain $\frac{r_0}{2} \leq r \leq r_0$.

(b) We graph $v = (0.5 - r)r^2$, and observe that the maximum indeed occurs at $v = \left(\frac{2}{3}\right)0.5 = \frac{1}{3}$.




$[0, 0.5]$ by $[-0.01, 0.03]$

61. The profit is given by

$$\begin{aligned} p(x) &= r(x) - c(x) \\ &= 6x - (x^3 - 6x^2 + 15x) \\ &= -x^3 + 6x^2 - 9x, \text{ for } x \geq 0. \end{aligned}$$

Then $p'(x) = -3x^2 + 12x - 9 = -3(x-1)(x-3)$, so the critical points occur at $x = 1$ and $x = 3$. Since $p'(x) < 0$ for $0 \leq x < 1$, $p'(x) > 0$ for $1 < x < 3$, and $p'(x) < 0$ for $x > 3$, the relative maxima occur at the endpoint $x = 0$ and at the critical point $x = 3$. Since $p(0) = p(3) = 0$, this means that for $x \geq 0$, the function $p(x)$ has its absolute maximum value at the points $(0, 0)$ and $(3, 0)$. This result can also be obtained graphically, as shown.



$[0, 5]$ by $[-8, 2]$

63. (a) According to the graph,
- $y'(0) = 0$
- .

(b) According to the graph, $y'(-L) = 0$.

(c) $y(0) = 0$, so $d = 0$.

Now $y'(x) = 3ax^2 + 2bx + c$, so $y'(0)$ implies that

$c = 0$. Therefore, $y(x) = ax^3 + bx^2$ and

$y'(x) = 3ax^2 + 2bx$. Then $y(-L) = -aL^3 + bL^2 = H$ and

$y'(-L) = 3aL^2 - 2bL = 0$, so we have two linear equations in the two unknowns a and b . The second

equation gives $b = \frac{3aL}{2}$. Substituting into the first

equation, we have $-aL^3 + \frac{3aL^3}{2} = H$, or

$\frac{aL^3}{2} = H$, so $a = 2\frac{H}{L^3}$. Therefore, $b = 3\frac{H}{L^2}$ and the

equation for y

is $y(x) = 2\frac{H}{L^3}x^3 + 3\frac{H}{L^2}x^2$, or

$$y(x) = H \left[2 \left(\frac{x}{L} \right)^3 + 3 \left(\frac{x}{L} \right)^2 \right].$$

65. (a) Let x_0 represent the fixed value of x at point P , so that P has coordinates (x_0, a) and let $m = f'(x_0)$ be the slope of line RT . Then the equation of line RT is $y = m(x - x_0) + a$. The y -intercept of this line is $m(0 - x_0) + a = a - mx_0$, and the x -intercept is the solution of $m(x - x_0) + a = 0$, or $x = \frac{mx_0 - a}{m}$. Let O designate the origin. Then (Area of triangle RST)

65. Continued

$$\begin{aligned}
 \text{(a)} &= 2 \text{ (Area of triangle } ORT) \\
 &= 2 \cdot \frac{1}{2} (x\text{-intercept of line } RT) (y\text{-intercept of line } RT) \\
 &= 2 \cdot \frac{1}{2} \left(\frac{mx_0 - a}{m} \right) (a - mx_0) \\
 &= -m \left(\frac{mx_0 - a}{m} \right) \left(\frac{mx_0 - a}{m} \right) \\
 &= - \left(\frac{mx_0 - a}{m} \right)^2 \\
 &= -m \left(x_0 - \frac{a}{m} \right)^2
 \end{aligned}$$

Substituting x for x_0 , $f'(x)$ for m , and $f(x)$ for a , we

$$\text{have } A(x) = -f'(x) \left[x - \frac{f(x)}{f'(x)} \right]^2.$$


(b) The domain is the open interval $(0, 10)$.

$$\text{To graph, let } y_1 = f(x) = 5 + 5\sqrt{1 - \frac{x^2}{100}},$$

$$y_2 = f'(x) = \text{NDER}(y_1), \text{ and}$$

$$y_3 = A(x) = -y_2 \left(x - \frac{y_1}{y_2} \right)^2.$$

The graph of the area function $y_3 = A(x)$ is shown below.



$[0, 10]$ by $[-100, 1000]$

The vertical asymptotes at $x = 0$ and $x = 10$ correspond to horizontal or vertical tangent lines, which do not form triangles.

(c) Using our expression for the y -intercept of the tangent line, the height of the triangle is

$$\begin{aligned}
 a - mx &= f(x) - f'(x) \cdot x \\
 &= 5 + \frac{1}{2}\sqrt{100 - x^2} - \frac{-x}{2\sqrt{100 - x^2}} x \\
 &= 5 + \frac{1}{2}\sqrt{100 - x^2} + \frac{x^2}{2\sqrt{100 - x^2}}
 \end{aligned}$$

We may use graphing methods or the analytic method in part (d) to find that the minimum value of $A(x)$ occurs at $x \approx 8.66$. Substituting this value into the expression above, the height of the triangle is 15. This is 3 times the y -coordinate of the center of the ellipse.

(d) Part (a) remains unchanged. The domain is $(0, C)$. To graph, note that

$$f(x) = B + B\sqrt{1 - \frac{x^2}{C^2}} = B + \frac{B}{C}\sqrt{C^2 - x^2} \text{ and}$$

$$f'(x) = \frac{B}{C} \frac{1}{2\sqrt{C^2 - x^2}} (-2x) = \frac{-Bx}{C\sqrt{C^2 - x^2}}.$$

Therefore, we have

$$\begin{aligned}
 A(x) &= -f'(x) \left[x - \frac{f(x)}{f'(x)} \right]^2 \\
 &= \frac{Bx}{C\sqrt{C^2 - x^2}} \left[x - \frac{B + \frac{B}{C}\sqrt{C^2 - x^2}}{\frac{-Bx}{C\sqrt{C^2 - x^2}}} \right]^2 \\
 &= \frac{Bx}{C\sqrt{C^2 - x^2}} \left[x - \frac{(BC + B\sqrt{C^2 - x^2})\sqrt{C^2 - x^2}}{-Bx} \right]^2 \\
 &= \frac{1}{BCx\sqrt{C^2 - x^2}} \left[Bx^2 + (BC + B\sqrt{C^2 - x^2})(\sqrt{C^2 - x^2}) \right]^2 \\
 &= \frac{1}{BCx\sqrt{C^2 - x^2}} \left[Bx^2 + BC\sqrt{C^2 - x^2} + B(C^2 - x^2) \right]^2 \\
 &= \frac{1}{BCx\sqrt{C^2 - x^2}} \left[BC(C + \sqrt{C^2 - x^2}) \right]^2 \\
 &= \frac{BC(C + \sqrt{C^2 - x^2})^2}{x\sqrt{C^2 - x^2}}
 \end{aligned}$$

$$\begin{aligned}
 A'(x) &= BC \cdot \frac{(x\sqrt{C^2 - x^2})(2)(C + \sqrt{C^2 - x^2}) \left(\frac{-x}{\sqrt{C^2 - x^2}} \right) - (C + \sqrt{C^2 - x^2})^2 \left(x \frac{-x}{\sqrt{C^2 - x^2}} + \sqrt{C^2 - x^2} (1) \right)}{x^2(C^2 - x^2)} \\
 &= \frac{BC(C + \sqrt{C^2 - x^2})}{x^2(C^2 - x^2)} \left[\frac{-2x^2 - (C + \sqrt{C^2 - x^2})}{\left(\frac{-x^2}{\sqrt{C^2 - x^2}} + \sqrt{C^2 - x^2} \right)} \right] \\
 &= \frac{BC(C + \sqrt{C^2 - x^2})}{x^2\sqrt{C^2 - x^2}} \left[\frac{-2x^2 + \frac{Cx^2}{\sqrt{C^2 - x^2}}}{-C\sqrt{C^2 - x^2} + x^2 - (C^2 - x^2)} \right] \\
 &= \frac{BC(C + \sqrt{C^2 - x^2})}{x^2(C^2 - x^2)} \left(\frac{Cx^2}{\sqrt{C^2 - x^2}} - C\sqrt{C^2 - x^2} - C^2 \right) \\
 &= \frac{BC(C + \sqrt{C^2 - x^2})}{x^2(C^2 - x^2)^{3/2}} \left[Cx^2 - C(C^2 - x^2) - C^2\sqrt{C^2 - x^2} \right] \\
 &= \frac{BC^2(C + \sqrt{C^2 - x^2})}{x^2(C^2 - x^2)^{3/2}} (2x^2 - C^2 - C\sqrt{C^2 - x^2})
 \end{aligned}$$

To find the critical points for $0 < x < C$, we solve:

$$\begin{aligned}
 2x^2 - C^2 &= C\sqrt{C^2 - x^2} \\
 4x^4 - 4C^2x^2 + C^4 &= C^4 = C^2x^2 \\
 4x^4 - 3C^2x^2 &= 0 \\
 x^2(4x^2 - 3C^2) &= 0
 \end{aligned}$$

The minimum value of $A(x)$ for $0 < x < C$ occurs at the

critical point $x = \frac{C\sqrt{3}}{2}$, or $x^2 = \frac{3C^2}{4}$. The corresponding triangle height is


65. Continued

$$\begin{aligned}
 a - mx &= f(x) - f'(x) \cdot x \\
 &= B + \frac{B}{C} \sqrt{C^2 - x^2} + \frac{Bx^2}{C\sqrt{C^2 - x^2}} \\
 &= B + \frac{B}{C} \sqrt{C^2 - \frac{3C^2}{4}} + \frac{B \left(\frac{3C^2}{4} \right)}{C\sqrt{C^2 - \frac{3C^2}{4}}} \\
 &= B + \frac{B}{C} \left(\frac{C}{2} \right) + \frac{3BC^2}{C^2} \\
 &= B + \frac{B}{2} + \frac{3B}{2} \\
 &= 3B
 \end{aligned}$$

This shows that the triangle has minimum area when its height is $3B$.

Section 4.5 Linearization and Newton's Method (pp. 233–245)

Exploration 1 Appreciating Local Linearity



$$y = (x^2 + 0.0001)^{1/4} + 0.9$$

The function appears to come to a point.

$$\begin{aligned}
 2. \quad f'(a) &= \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} \\
 &= \lim_{x \rightarrow a} \frac{(x^2 + 0.0001)^{1/4} + 0.9 - ((0 + 0.0001)^{1/4} + 0.9)}{x - 0} \\
 &= \lim_{x \rightarrow a} \frac{(x^2 + 0.0001)^{1/4} - 0.1}{x} = 0
 \end{aligned}$$

$f(x)$ is differentiable at $x = 0$, and the equation of the tangent line is $y = 1$.

3. The graph of the function at that point seems to become the graph of a straight line with repeated zooming.

4. The graph will eventually look like the tangent line.


Exploration 2 Using Newton's Method on Your Calculator

See text page 237.

Quick Review 4.5


$$1. \quad \frac{dy}{dx} = \cos(x^2 + 1) \cdot \frac{d}{dx}(x^2 + 1) = 2x \cos(x^2 + 1)$$

$$\begin{aligned}
 2. \quad \frac{dy}{dx} &= \frac{(x+1)(1-\sin x) - (x+\cos x)(1)}{(x+1)^2} \\
 &= \frac{x - x \sin x + 1 - \sin x - x - \cos x}{(x+1)^2} \\
 &= \frac{1 - \cos x - (x+1) \sin x}{(x+1)^2}
 \end{aligned}$$



$$[-2, 6] \text{ by } [-3, 3]$$

$$x \approx -0.567$$



$$[-4, 4] \text{ by } [-10, 10]$$

$$x \approx -0.322$$

$$5. \quad f'(x) = (x)(-e^{-x}) + (e^{-x})(1) = e^{-x} - xe^{-x}$$

$$f'(0) = 1$$

The line passes through $(0, 1)$ and has slope 1. Its equation is $y = x + 1$.

$$6. \quad f'(x) = (x)(-e^{-x}) + (e^{-x})(1) = e^{-x} - xe^{-x}$$

$$f'(-1) = e^1 - (-e^1) = 2e$$

The line passes through $(-1, -e + 1)$ and has slope $2e$. Its equation is $y = 2e(x + 1) + (-e + 1)$, or $y = 2ex + e + 1$.

$$7. \quad \text{(a) } x + 1 = 0$$

$$x = -1$$

$$\text{(b) } 2ex + e + 1 = 0$$

$$2ex = -(e + 1)$$

$$x = -\frac{e + 1}{2e} \approx -0.684$$

8. $f'(x) = 3x^2 - 4$

$f'(1) = 3(1)^2 - 4 = -1$

Since $f(1) = -2$ and $f'(1) = -1$, the graph of $g(x)$ passes through $(1, -2)$ and has slope -1 . Its equation is

$g(x) = -1(x-1) + (-2), \text{ or } g(x) = -x - 1.$


| x | $f(x)$ | $g(x)$ |
|-----|--------|--------|
| 0.7 | -1.457 | -1.7 |
| 0.8 | -1.688 | -1.8 |
| 0.9 | -1.871 | -1.9 |
| 1.0 | -2 | -2 |
| 1.1 | -2.069 | -2.1 |
| 1.2 | -2.072 | -2.2 |
| 1.3 | -2.003 | -2.3 |

9. $f'(x) = \cos x$

$f'(1.5) = \cos 1.5$

Since $f(1.5) = \sin 1.5$ and $f'(1.5) = \cos 1.5$, the tangent line passes through $(1.5, \sin 1.5)$ and has slope $\cos 1.5$. Its equation is $y = (\cos 1.5)(x - 1.5) + \sin 1.5$, or approximately

$y = 0.071x + 0.891$




$[0, \pi]$ by $[-0.2, 1.3]$

10. For $x > 3$, $f'(x) = \frac{1}{2\sqrt{x-3}}$, and so $f'(4) = \frac{1}{2}$. Since

$f(4) = 1$ and $f'(4) = \frac{1}{2}$, the tangent line passes through

$(4, 1)$ and has slope $\frac{1}{2}$. Its equation is

$y = \frac{1}{2}(x-4) + 1, \text{ or } y = \frac{1}{2}x - 1.$



$[-1, 7]$ by $[-2, 2]$

Section 4.5 Exercises

1. (a) $f'(x) = 3x^2 - 2$

We have $f(2) = 7$ and $f'(2) = 10$.

$$\begin{aligned} L(x) &= f(2) + f'(2)(x-2) \\ &= 7 + 10(x-2) \\ &= 10x - 13 \end{aligned}$$

(b) Since $f(2.1) = 8.061$ and $L(2.1) = 8$, the approximation differs from the true value in absolute value by less than 10^{-1} .

2. (a) $f'(x) = \frac{1}{2\sqrt{x^2+9}}(2x) = \frac{x}{\sqrt{x^2+9}}$

We have $f(-4) = 5$ and $f'(-4) = -\frac{4}{5}$.

$$\begin{aligned} L(x) &= f(-4) + f'(-4)(x - (-4)) \\ &= 5 - \frac{4}{5}(x+4) \\ &= -\frac{4}{5}x + \frac{9}{5} \end{aligned}$$

(b) Since $f(-3.9) \approx 4.9204$ and $L(-3.9) = 4.92$, the approximation differs from the true value by less than 10^{-3} .

3. (a) $f'(x) = 1 - x^{-2}$

We have $f(1) = 2$ and $f'(1) = 0$.

$$\begin{aligned} L(x) &= f(1) + f'(1)(x-1) \\ &= 2 + 0(x-1) \\ &= 2 \end{aligned}$$

(b) Since $f(1.1) \approx 2.009$ and $L(1.1) = 2$, the approximation differs from the true value by less than 10^{-2} .

4. (a) $f'(x) = \frac{1}{x+1}$

We have $f(0) = 0$ and $f'(0) = 1$.

$$\begin{aligned} L(x) &= f(0) + f'(0)(x-0) \\ &= 0 + 1x \\ &= x \end{aligned}$$

(b) Since $f(0.1) \approx 0.0953$ and $L(0.1) = 0.1$ the approximation differs from the true value by less than 10^{-2} .

5. (a) $f'(x) = \sec^2 x$

We have $f(\pi) = 0$ and $f'(\pi) = 1$.

$$\begin{aligned} L(x) &= f(\pi) + f'(\pi)(x-\pi) \\ &= 0 + 1(x-\pi) \\ &= x - \pi \end{aligned}$$

(b) Since $f(\pi+0.1) \approx 0.10033$ and $L(\pi+0.1) = 0.1$, the approximation differs from the true value in absolute value by less than 10^{-3} .

6. (a) $f'(x) = -\frac{1}{\sqrt{1-x^2}}$

We have $f(0) = \frac{\pi}{2}$ and $f'(0) = -1$.

$$\begin{aligned} L(x) &= f(0) + f'(0)(x-0) \\ &= \frac{\pi}{2} + (-1)(x-0) \\ &= -x + \frac{\pi}{2} \end{aligned}$$

6. Continued

(b) Since $f(0.1) \approx 1.47063$ and $L(0.1) \approx 1.47080$, the approximation differs from the true value in absolute value by less than 10^{-3} .

$$7. f'(x) = k(1+x)^{k-1}$$

We have $f(0) = 1$ and $f'(0) = k$.

$$\begin{aligned} L(x) &= f(0) + f'(0)(x-0) \\ &= 1 + k(x-0) \\ &= 1 + kx \end{aligned}$$

$$8. (a) (1.002)^{100} = (1+0.002)^{100} \approx 1 + (100)(0.002) = 1.2;$$

$$|1.002^{100} - 1.2| \approx 0.021 < 10^{-1}$$

$$(b) \sqrt[3]{1.009} = (1+0.009)^{1/3} \approx 1 + \frac{1}{3}(0.009) = 1.003;$$

$$|\sqrt[3]{1.009} - 1.003| \approx 9 \times 10^{-6} < 10^{-5}$$

$$9. (a) f(x) = (1-x)^6 = [1+(-x)]^6 \approx 1 + 6(-x) = 1 - 6x$$

$$(b) f(x) = \frac{2}{1-x} = 2[1+(-x)]^{-1} \approx 2[1+(-1)(-x)] = 2 + 2x$$

$$(c) f(x) = (1+x)^{-1/2} \approx 1 + \left(-\frac{1}{2}\right)x = 1 - \frac{x}{2}$$

$$10. (a) f(x) = (4+3x)^{1/3} = 4^{1/3} \left(1 + \frac{3x}{4}\right)^{1/3} \\ \approx 4^{1/3} \left(1 + \frac{1}{3} \left(\frac{3x}{4}\right)\right) = 4^{1/3} \left(1 + \frac{x}{4}\right)$$

$$(b) f(x) = \sqrt{2+x^2} = \sqrt{2} \left(1 + \frac{x^2}{2}\right)^{1/2} \\ \approx \sqrt{2} \left(1 + \frac{1}{2} \left(\frac{x^2}{2}\right)\right) = \sqrt{2} \left(1 + \frac{x^2}{4}\right)$$

$$(c) f(x) = \left(1 - \frac{1}{2+x}\right)^{2/3} = \left[1 + \left(-\frac{1}{2+x}\right)\right]^{2/3} \\ \approx 1 + \frac{2}{3} \left(-\frac{1}{2+x}\right) = 1 - \frac{2}{6+3x}$$

$$11. x = 100$$

$$f'(100) = \frac{1}{2}(100)^{-1/2} = 0.05$$

$$f(100) = 10 + 0.05(101 - 100) = 10.05$$

$$12. x = 27$$

$$f'(27) = \frac{1}{3}(27)^{-2/3} = \frac{1}{27}$$

$$f(27) = 3 + (1/27)(26 - 27)$$

$$y = 3 - \frac{1}{27} \approx 2.962$$

$$13. x = 1000$$

$$f'(1000) = \frac{1}{3}(1000)^{-2/3} = \frac{1}{300}$$

$$y = 10 + (1/300)(x - 1000)$$

$$y = 10 - \frac{1}{150} = 9.99\bar{3}$$

$$14. x = 81$$

$$f'(81) = \frac{1}{2}(81)^{-1/2} = \frac{1}{18}$$

$$y = 9 + \frac{1}{18}(80 - 81)$$

$$y = 9 - \frac{1}{18} = 8.9\bar{4}$$

15. Let $f(x) = x^3 + x - 1$. Then $f'(x) = 3x^2 + 1$ and

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{x_n^3 + x_n - 1}{3x_n^2 + 1}$$

Note that f is cubic and f' is always positive, so there is exactly one solution. We choose $x_1 = 0$.

$$x_1 = 0$$

$$x_2 = 1$$

$$x_3 = 0.75$$

$$x_4 \approx 0.6860465$$

$$x_5 \approx 0.6823396$$

$$x_6 \approx 0.6823278$$


$$x_7 \approx 0.6823278$$

Solution: $x \approx 0.682328$.

16. Let $f(x) = x^4 + x - 3$. Then $f'(x) = 4x^3 + 1$ and

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{x_n^4 + x_n - 3}{4x_n^3 + 1}$$

The graph of $y = f(x)$ shows that $f(x) = 0$ has two solutions.



$[-3, 3]$ by $[-4, 4]$

$$x_1 = -1.5$$

$$x_1 = 1.2$$

$$x_2 = -1.455$$

$$x_2 \approx 1.6541962$$

$$x_3 = -1.4526332$$

$$x_3 \approx 1.1640373$$

$$x_4 = -1.4526269$$

$$x_4 \approx 1.1640351$$

$$x_5 = -1.4526269$$

$$x_5 \approx 1.1640351$$

Solution: $x \approx -1.452627, 1.164035$


17. Let $f(x) = x^2 - 2x + 1 - \sin x$.

Then $f'(x) = 2x - 2\cos x$ and

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{x_n^2 - 2x_n + 1 - \sin x_n}{2x_n - 2\cos x_n}$$

17. Continued

The graph of $y = f(x)$ shows that $f(x) = 0$ has two solutions



$[-4, 4]$ by $[-3, 3]$

$$\begin{array}{ll} x_1 = 0.3 & x_1 = 2 \\ x_2 \approx 0.3825699 & x_2 \approx 1.9624598 \\ x_3 \approx 0.3862295 & x_3 \approx 1.9615695 \\ x_4 \approx 0.3862369 & x_4 \approx 1.9615690 \\ x_5 \approx 0.3862369 & x_5 \approx 1.9615690 \end{array}$$

Solutions: $x \approx 0.386237, 1.961569$

18. Let $f(x) = x^4 - 2$. Then $f'(x) = 4x^3$ and

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{x_n^4 - 2}{4x_n^3}$$

Note that $f(x) = 0$ clearly has two solutions, namely

$x = \pm\sqrt[4]{2}$. We use Newton's method to find the decimal equivalents.

$$\begin{array}{l} x_1 = 1.5 \\ x_2 \approx 1.2731481 \\ x_3 \approx 1.1971498 \\ x_4 \approx 1.1892858 \\ x_5 \approx 1.1892071 \\ x_6 \approx 1.1892071 \end{array}$$

Solutions: $x \approx \pm 1.189207$

19. (a) Since $\frac{dy}{dx} = 3x^2 - 3$, $dy = (3x^2 - 3)dx$.

(b) At the given values,

$$dy = (3 \cdot 2^2 - 3)(0.05) = 9(0.05) = 0.45.$$

20. (a) Since $\frac{dy}{dx} = \frac{(1+x^2)(2) - (2x)(2x)}{(1+x^2)^2} = \frac{2-2x^2}{(1+x^2)^2}$,

$$dy = \frac{2-2x^2}{(1+x^2)^2} dx.$$

(b) At the given values,

$$\begin{aligned} dy &= \frac{2-2(-2)^2}{[1+(-2)^2]^2} (0.1) = \frac{2-8}{5^2} (0.1) \\ &= -0.024. \end{aligned}$$

21. (a) Since $\frac{dy}{dx} = (x^2) \left(\frac{1}{x} \right) + (\ln x)(2x) = 2x \ln x + x$,

$$dy = (2x \ln x + x) dx.$$

(b) At the given values,

$$dy = [2(1) \ln(1) + 1](0.01) = 1(0.01) = 0.01$$

$$\begin{aligned} 22. \text{ (a) Since } \frac{dy}{dx} &= (x) \left(\frac{1}{2\sqrt{1-x^2}} \right) (-2x) + (\sqrt{1-x^2})(1) \\ &= \frac{-x^2}{\sqrt{1-x^2}} + \sqrt{1-x^2} = \frac{-x^2 + (1-x^2)}{\sqrt{1-x^2}} = \frac{1-2x^2}{\sqrt{1-x^2}}, \\ dy &= \frac{1-2x^2}{\sqrt{1-x^2}} dx. \end{aligned}$$

$$\text{(b) At the given values, } dy = \frac{1-2(0)^2}{\sqrt{1-(0)^2}} (-0.2) = -0.2.$$

23. (a) Since $\frac{dy}{dx} = e^{\sin x} \cos x$, $dy = (\cos x)e^{\sin x} dx$.

(b) At the given values,

$$dy = (\cos \pi)(e^{\sin \pi})(-0.1) = (-1)(1)(-0.1) = 0.1.$$

$$\begin{aligned} 24. \text{ (a) Since } \frac{dy}{dx} &= -3 \csc \left(1 - \frac{x}{3} \right) \cot \left(1 - \frac{x}{3} \right) \left(-\frac{1}{3} \right) \\ &= \csc \left(1 - \frac{x}{3} \right) \cot \left(1 - \frac{x}{3} \right), \\ dy &= \csc \left(1 - \frac{x}{3} \right) \cot \left(1 - \frac{x}{3} \right) dx. \end{aligned}$$

(b) At the given values,

$$\begin{aligned} dy &= \csc \left(1 - \frac{1}{3} \right) \cot \left(1 - \frac{1}{3} \right) (0.1) \\ &= 0.1 \csc \frac{2}{3} \cot \frac{2}{3} \approx 0.205525 \end{aligned}$$

25. (a) $y + xy - x = 0$

$$y(1+x) = x$$

$$y = \frac{x}{x+1}$$

$$\text{Since } \frac{dy}{dx} = \frac{(x+1)(1) - (x)(1)}{(x+1)^2} = \frac{1}{(x+1)^2},$$

$$dy = \frac{dx}{(x+1)^2}.$$

(b) At the given values,

$$dy = \frac{0.01}{(0+1)^2} = 0.01.$$

26. (a) $2y = x^2 - xy$

$$2dy = 2x dx - x dy - y dx$$

$$dy(2+x) = (2x-y) dx$$

$$dy = \left(\frac{2x-y}{2+x} \right) dx$$

(b) At the given values, and $y = 1$ from the original

$$\text{equation, } dy = \left(\frac{2(2)-1}{2+2} \right) (-0.05) = -0.0375$$

$$27. \frac{dy}{dx} = \sqrt{1-x^2}$$

$$dy = \left(-\frac{2x}{2\sqrt{1-x^2}} \right) dx$$

$$dy = -\frac{x}{\sqrt{1-x^2}} dx$$

$$28. \frac{dy}{dx} = e^{5x} + x^5$$

$$dy = (5e^{5x} + 5x^4) dx$$

$$29. \frac{dy}{dx} = \tan^{-1} 4x$$

$$\frac{d}{dx} \tan^{-1} u = \frac{1}{1+u^2} \frac{du}{dx}$$

$$u = 4x$$

$$\frac{du}{dx} = 4$$

$$dy = \left(\frac{4}{1+16x^2} \right) dx$$

$$30. \frac{dy}{dx} = (8^x + x^8)$$

$$\frac{d}{dx} a^x = (\ln a) a^x$$

$$dy = (8^x \ln 8 + 8x^7) dx$$

$$31. (a) \Delta f = f(0.1) - f(0) = 0.21 - 0 = 0.21$$

$$(b) \text{ Since } f'(x) = 2x + 2, f'(0) = 2.$$

$$\text{Therefore, } df = 2 dx = 2(0.1) = 0.2.$$

$$(c) |\Delta f - df| = |0.21 - 0.2| = 0.01$$

$$32. (a) \Delta f = f(1.1) - f(1) = 0.231 - 0 = 0.231$$

$$(b) \text{ Since } f'(x) = 3x^2 - 1, f'(1) = 2.$$

$$\text{Therefore, } df = 2 dx = 2(0.1) = 0.2.$$

$$(c) |\Delta f - df| = |0.231 - 0.2| = 0.031$$

$$33. (a) \Delta f = f(0.55) - f(0.5) = \frac{20}{11} - 2 = -\frac{2}{11}$$

$$(b) \text{ Since } f'(x) = -x^{-2}, f'(0.5) = -4.$$

$$\text{Therefore, } df = -4 dx = -4(0.05) = -0.2 = -\frac{1}{5}$$

$$(c) |\Delta f - df| = \left| -\frac{2}{11} + \frac{1}{5} \right| = \frac{1}{55}$$

$$34. (a) \Delta f = f(1.01) - f(1) = 1.04060401 - 1 = 0.04060401$$

$$(b) \text{ Since } f'(x) = 4x^3, f'(1) = 4.$$

$$\text{Therefore, } df = 4 dx = 4(0.01) = 0.04.$$

$$(c) |\Delta f - df| = |0.04060401 - 0.04| = 0.00060401$$

$$35. \text{ Note that } \frac{dV}{dr} = 4\pi r^2, dV = 4\pi r^2 dr. \text{ When } r \text{ changes from } a \text{ to } a + dr, \text{ the change in volume is approximately } 4\pi a^2 dr.$$

$$36. \text{ Note that } \frac{dS}{dr} = 8\pi r, \text{ so } dS = 8\pi r dr. \text{ When } r \text{ changes from } a \text{ to } a + dr, \text{ the change in surface area is approximately } 8\pi a dr.$$

$$37. \text{ Note that } \frac{dV}{dx} = 3x^2, \text{ so } dV = 3x^2 dx. \text{ When } x \text{ changes from } a \text{ to } a + dx, \text{ the change in volume is approximately } 3a^2 dx.$$

$$38. \text{ Note that } \frac{dS}{dx} = 12x, \text{ so } dS = 12x dx. \text{ When } x \text{ changes from } a \text{ to } a + dx, \text{ the change in surface area is approximately } 12a dx.$$

$$39. \text{ Note that } \frac{dV}{dr} = 2\pi rh, \text{ so } dV = 2\pi rh dr. \text{ When } r \text{ changes from } a \text{ to } a + dr, \text{ the change in volume is approximately}$$

$$40. \text{ Note that } \frac{dS}{dh} = 2\pi r, \text{ so } dS = 2\pi r dh. \text{ When } h \text{ changes from } a \text{ to } a + dh, \text{ the change in lateral surface area is approximately } 2\pi r dh.$$

$$41. A = \pi r^2 \\ dA = 2\pi r dr \\ dA = 2\pi(10)(0.1) = 6.3 \text{ in}^2$$

$$42. v = \frac{4}{3}\pi r^3 \\ dV = 4\pi r^2 dr \\ dV = 4\pi(8)^2(0.3) = 241 \text{ in}^2$$

$$43. v = s^3 \\ dV = 3s^2 ds \\ dV = 3(15)^2(0.2) = 135 \text{ cm}^2$$

$$44. A = \frac{\sqrt{3}}{4}s^2 \\ dA = \frac{\sqrt{3}}{2}s ds \\ dA = \frac{\sqrt{3}}{2}(20)(0.5) = 8.7 \text{ cm}^2$$

$$45. (a) \text{ Note that } f'(0) = \cos 0 = 1. \\ L(x) = f(0) + f'(0)(x-0) = 1 + 1x = x + 1$$

$$(b) f(0.1) \approx L(0.1) = 1.1$$

45. Continued

(c) The actual value is less than 1.1. This is because the derivative is decreasing over the interval $[0, 0.1]$, which means that the graph of $f(x)$ is concave down and lies below its linearization in this interval.

46. (a) Note that $A = \pi r^2$ and $\frac{dA}{dr} = 2\pi r$, so $dA = 2\pi r dr$.

When r changes from a to $a + dr$, the change in area is approximately $2\pi a dr$. Substituting 2 for a and 0.02 for dr , the change in area is approximately $2\pi(2)(0.02) = 0.08\pi \approx 0.2513$

$$(b) \frac{dA}{A} = \frac{0.08\pi}{4\pi} = 0.02 = 2\%$$

47. Let A = cross section area, C = circumference, and

$$D = \text{diameter. Then } D = \frac{C}{\pi}, \text{ so } \frac{dD}{dC} = \frac{1}{\pi}$$

$$\text{and } dD = \frac{1}{\pi} dC. \text{ Also, } A = \pi \left(\frac{D}{2}\right)^2 = \pi \left(\frac{C}{2\pi}\right)^2 = \frac{C^2}{4\pi},$$

so $\frac{dA}{dC} = \frac{C}{2\pi}$ and $dA = \frac{C}{2\pi} dC$. When C increases from 10π in. to 10π + 2 in. the diameter increases by

$$dD = \frac{1}{\pi}(2) = \frac{2}{\pi} \approx 0.6366 \text{ in. and the area increases by}$$

$$\text{approximately } dA = \frac{10\pi}{2\pi}(2) = 10 \text{ in}^2.$$

48. Let x = edge length and V = volume. Then $V = x^3$, and

so $dV = 3x^2 dx$. With $x = 10$ cm and $dx = 0.01x = 0.1$ cm, we have $V = 10^3 = 1000 \text{ cm}^3$ and

$dV = 3(10)^2(0.1) = 30 \text{ cm}^3$, so the percentage error in the volume measurement is approximately

$$\frac{dV}{V} = \frac{30}{1000} = 0.03 = 3\%.$$

49. Let x = side length and A = area. Then $A = x^2$ and

$$\frac{dA}{dx} = 2x, \text{ so } dA = 2x dx. \text{ We want } |dA| \leq 0.02A, \text{ which}$$

gives $|2x dx| \leq 0.02x^2$, or $|dx| \leq 0.01x$. The side length should be measured with an error of no more than 1%.

For $\theta = 75^\circ = \frac{5\pi}{12}$ radians, we have

$$|d\theta| < 0.04 \sin \frac{5\pi}{12} \cos \frac{5\pi}{12} = 0.01 \text{ radian. The angle should be}$$

measured with an error of less than 0.01 radian (or approximately 0.57 degrees), which is a percentage error of approximately 0.76%.

50. (a) Note that $V = \pi r^2 h = 10\pi r^2 = 2.5\pi D^2$, where D is the interior diameter of the tank. Then $\frac{dV}{dD} = 5\pi D$,

so $dV = 5\pi D dD$. We want $|dV| \leq 0.01V$, which

gives $|5\pi D dD| \leq 0.01(2.5\pi D^2)$, or $|dD| \leq 0.005D$. The interior diameter should be measured with an error of no more than 0.5%.

(b) Now we let D represent the exterior diameter of the tank, and we assume that the paint coverage rate (number of square feet covered per gallon of paint) is known precisely. Then, to determine the amount of paint within 5%, we need to calculate the lateral surface area S with an error of no more than 5%. Note that

$$S = 2\pi r h = 10\pi D, \text{ so } \frac{dS}{dD} = 10\pi \text{ and } dS = 10\pi dD. \text{ We}$$

want $|dS| \leq 0.05S$, which gives $|10\pi dD| \leq 0.05(10\pi D)$, or $dD \leq 0.05D$. The exterior diameter should be measured with an error of no more than 5%.

51. Note that $V = \pi r^2 h$, where h is constant. Then $\frac{dV}{dr} = 2\pi r h$.

The percent change is given by

$$\frac{dV}{V} = \frac{2\pi r h dr}{\pi r^2 h} = 2 \frac{dr}{r} = 2 \frac{0.1\%}{r} = 0.2\%.$$

52. Note that $\frac{dV}{dh} = 3\pi h^2$, so $dV = 3\pi h^2 dh$. We want

$$|dV| \leq 0.01V, \text{ which gives } |3\pi h^2 dh| \leq 0.01(\pi h^3),$$

or $|dh| \leq \frac{0.01h}{3}$. The height should be measured with an

error of no more than $\frac{1}{3}\%$.

53. Since $V = \frac{4}{3}\pi r^3$, we have

$$dV = 4\pi r^2 dr = 4\pi r^2 \left(\frac{1}{16\pi}\right) = \frac{r^2}{4}. \text{ The volume error in}$$

each case is simply $\frac{r^2}{4} \text{ in}^3$.

| Sphere Type | True Radius | Tape error | Radius Error | Volume Error |
|-------------|-------------|------------|--------------|------------------------|
| Orange | 2 in. | 1/8 in. | 1/16π in. | 1 in. ³ |
| Melon | 4 in. | 1/8 in. | 1/16π in. | 4 in. ³ |
| Beach Ball | 7 in. | 1/8 in. | 1/16π in. | 12.25 in. ³ |

54. Since $A = 4\pi r^2$, we have $dA = 8\pi r dr = 8\pi r \left(\frac{1}{16\pi}\right) = \frac{r}{2}$.

The surface area error in each case is simply $\frac{r}{2} \text{ in.}^2$.

| Sphere Type | True Radius | Tape Error | Radius Error | Volume Error |
|-------------|-------------|------------|--------------|----------------------|
| Orange | 2 in. | 1/8 in. | 1/16π in. | 1 in. ² |
| Melon | 4 in. | 1/8 in. | 1/16π in. | 2 in. ² |
| Beach Ball | 7 in. | 1/8 in. | 1/16π in. | 3.5 in. ² |

55. We have $\frac{dW}{dg} = -bg^{-2}$, so $dW = -bg^{-2}dg$.

Then $\frac{dW_{\text{moon}}}{dW_{\text{earth}}} = \frac{-b(5.2)^{-2}dg}{-b(32)^{-2}dg} = \frac{32^2}{5.2^2} \approx 37.87$. The ratio is about 37.87 to 1.

56. (a) Note that $T = 2\pi L^{1/2}g^{-1/2}$, so $\frac{dT}{dg} = -\pi L^{1/2}g^{-3/2}$ and

$$dT = -\pi L^{1/2}g^{-3/2}dg.$$

(b) Note that dT and dg have opposite signs. Thus, if g increases, T decreases and the clock speeds up.

(c)
$$-\pi L^{1/2}g^{-3/2}dg = dT$$

$$-\pi(100)^{1/2}(980)^{-3/2}dg = 0.001$$

$$dg \approx -0.9765$$

Since $dg = -0.9765$, $g = 980 - 0.9765 = 979.0235$.

57. True. A look at the graph reveals the problem. The graph decreases after $x=1$ toward a horizontal asymptote of $x=0$, so the x -intercepts of the tangent lines keep getting bigger without approaching a zero.

58. False. By the product rule, $d(uv) = udv + vdu$.

59. B. $f(x) = e^x$
 $f'(x) = e^x$
 $L(x) = e^1 + e^1(x-1)$
 $L(x) = ex$

60. A. $y = \tan x$
 $dy = (\sec^2 x)dx = (\sec^2 \pi)0.5$
 $dy = -0.25$

61. D. $f(x) = x - x^3 + 2$
 $f'(x) = 1 - 3x^2$

$$x_{n+1} = x_n - \frac{x_n x_n^3 + 2}{1 - 3x_n^2}$$

$$x_2 = 1 - \frac{1 - (1)^3 + 2}{1 - 3(1)^2} = 2$$

$$x_3 = 2 - \frac{2 - (2)^3 + 2}{1 - 3(2)^2} = \frac{18}{11}$$

62. A. $f(x) = \sqrt[3]{x}$, $x = 64$

$$f'(64) = \frac{1}{3}(64)^{-2/3} = \frac{1}{48}$$

$$\sqrt[3]{66} = 4 + \frac{1}{48}(66 - 64)$$

$$\sqrt[3]{66} = 4.042$$

The calculator returns 4.041, or a 0.01% difference.


63. If $f'(x) \neq 0$, we have $x_2 = x_1 - \frac{f(x_1)}{f'(x_1)} = x_1 - \frac{0}{f'(x_1)} = x_1$.

Therefore, $x_2 = x_1$, and all later approximations are also equal to x_1 .

64. If $x_1 = h$, then $f'(x_1) = \frac{1}{2h^{1/2}}$ and

$$x_2 = h - \frac{h^{1/2}}{\frac{1}{2h^{1/2}}} = h - 2h = -h. \text{ If } x_1 = -h, \text{ then}$$

$$f'(x_1) = -\frac{1}{2\sqrt{h}} \text{ and } x_2 = -h - \frac{h^{1/2}}{-\frac{1}{2h^{1/2}}} = -h + 2h = h$$




$[-3, 3]$ by $[-0.5, 2]$

65. Note that $f'(x) = \frac{1}{3}x^{-2/3}$ and so

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{x_n^{1/3}}{\frac{x_n^{-2/3}}{3}} = x_n - 3x_n = -2x_n. \text{ For}$$

$$x_1 = 1, \text{ we have } x_2 = -2, x_3 = 4, x_4 = -8, \text{ and}$$

$$x_5 = 16; |x_n| = 2^{n-1}.$$



$[-10, 10]$ by $[-3, 3]$

66. (a) i. $Q(a) = f(a)$ implies that $b_0 = f(a)$.

ii. Since $Q'(x) = b_1 + 2b_2(x - a)$, $Q'(a) = f'(a)$ implies that $b_1 = f'(a)$.

66. Continued

iii. Since $Q''(x) = 2b_2$, $Q''(a) = f''(a)$ implies that

$$b_2 = \frac{f''(a)}{2}$$


In summary, $b_0 = f(a)$, $b_1 = f'(a)$, and $b_2 = \frac{f''(a)}{2}$.

(b) $f(x) = (1-x)^{-1}$
 $f'(x) = -1(1-x)^{-2}(-1) = (1-x)^{-2}$
 $f''(x) = -2(1-x)^{-3}(-1) = 2(1-x)^{-3}$

Since $f(0) = 1$, $f'(0) = 1$, and $f''(0) = 2$, the coefficients are

$$b_0 = 1, b_1 = 1, \text{ and } b_2 = \frac{2}{2} = 1. \text{ The quadratic approximation}$$

is $Q(x) = 1 + x + x^2$.




$[-2.35, 2.35]$ by $[-1.25, 3.25]$

As one zooms in, the two graphs quickly become indistinguishable. They appear to be identical.

(d) $g(x) = x^{-1}$
 $g'(x) = -x^{-2}$
 $g''(x) = 2x^{-3}$
 Since $g(1) = 1$, $g'(1) = -1$, and $g''(1) = 2$, the coefficients are $b_0 = 1$, $b_1 = -1$, and $b_2 = \frac{2}{2} = 1$. The

quadratic approximation is $Q(x) = 1 - (x-1) + (x-1)^2$.



$[-1.35, 3.35]$ by $[-1.25, 3.25]$


As one zooms in, the two graphs quickly become indistinguishable. They appear to be identical.

(e) $h(x) = (1+x)^{1/2}$
 $h'(x) = \frac{1}{2}(1+x)^{-1/2}$
 $h''(x) = -\frac{1}{4}(1+x)^{-3/2}$

Since $h(0) = 1$, $h'(0) = \frac{1}{2}$, and $h''(0) = -\frac{1}{4}$, the

coefficients are $b_0 = 1$, $b_1 = \frac{1}{2}$, and $b_2 = \frac{-\frac{1}{4}}{2} = -\frac{1}{8}$.

The quadratic approximation is $Q(x) = 1 + \frac{x}{2} - \frac{x^2}{8}$.



$[-1.35, 3.35]$ by $[-1.25, 3.25]$

As one zooms in, the two graphs quickly become indistinguishable. They appear to be identical.

(f) The linearization of any differentiable function $u(x)$ at $x = a$ is $L(x) = u(a) + u'(a)(x-a) = b_0 + b_1(x-a)$, where b_0 and b_1 are the coefficients of the constant and linear terms of the quadratic approximation. Thus, the linearization for $f(x)$ at $x = 0$ is $1 + x$; the linearization for $g(x)$ at $x = 1$ is $1 - (x-1)$ or $2 - x$; and the linearization for $h(x)$ at $x = 0$ is $1 + \frac{x}{2}$.

67. Finding a zero of $\sin x$ by Newton's method would use the

recursive formula $x_{n+1} = x_n - \frac{\sin(x_n)}{\cos(x_n)} = x_n - \tan x_n$, and that

is exactly what the calculator would be doing. Any zero of $\sin x$ would be a multiple of π .

68. Just multiply the corresponding derivative formulas by dx .

(a) Since $\frac{d}{dx}(c) = 0$, $d(c) = 0$.

(b) Since $\frac{d}{dx}(cu) = c \frac{du}{dx}$, $d(cu) = c du$.

(c) Since $\frac{d}{dx}(u+v) = \frac{du}{dx} + \frac{dv}{dx}$, $d(u+v) = du + dv$

(d) Since $\frac{d}{dx}(u \cdot v) = u \frac{dv}{dx} + v \frac{du}{dx}$, $d(u \cdot v) = u dv + v du$.

(e) Since $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$, $d\left(\frac{u}{v}\right) = \frac{v du - u dv}{v^2}$.

(f) Since $\frac{d}{dx}u^n = nu^{n-1} \frac{du}{dx}$, $d(u^n) = nu^{n-1} du$.

69. $\lim_{x \rightarrow 0} \frac{\tan x}{x} = \lim_{x \rightarrow 0} \frac{\sin x / \cos x}{x}$
 $= \lim_{x \rightarrow 0} \left(\frac{1}{\cos x} \cdot \frac{\sin x}{x} \right)$
 $= \left(\lim_{x \rightarrow 0} \frac{1}{\cos x} \right) \left(\lim_{x \rightarrow 0} \frac{\sin x}{x} \right)$
 $= (1)(1) = 1.$

70. $g(a) = c$, so if $E(a) = 0$, then $g(a) = f(a)$ and $c = f(a)$.

Then $E(x) = f(x) - g(x) = f(x) - f(a) - m(x - a)$.

$$\text{Thus, } \frac{E(x)}{x-a} = \frac{f(x) - f(a)}{x-a} - m.$$

$$\lim_{x \rightarrow a} \frac{f(x) - f(a)}{x-a} = f'(a), \text{ so } \lim_{x \rightarrow a} \frac{E(x)}{x-a} = f'(a) - m.$$

Therefore, if the limit of $\frac{E(x)}{x-a}$ is zero, then $m = f'(a)$ and

$$g(x) = L(x).$$

$$71. f'(x) = \frac{1}{2\sqrt{x+1}} + \cos x$$

$$\text{We have } f(0) = 1 \text{ and } f'(0) = \frac{3}{2}$$

$$L(x) = f(0) + f'(0)(x-0)$$

$$= 1 + \frac{3}{2}x$$

The linearization is the sum of the two individual

linearizations, which are x for $\sin x$ and $1 + \frac{1}{2}x$ for $\sqrt{x+1}$.

Section 4.6 Related Rates (pp. 246–255)

Exploration 1 Sliding Ladder

1. Here the x -axis represents the ground and the y -axis represents the wall. The curve (x_1, y_1) gives the position of the bottom of the ladder (distance from the wall) at any time t in $0 \leq t \leq 5$. The curve (x_2, y_2) gives the position of the top of the ladder at any time in $0 \leq t \leq 5$.

2. $0 \leq t \leq 5$

4. This is a snapshot at $t \approx 3$. 1. The top of the ladder is moving down the y -axis and the bottom of the ladder is moving to the right on the x -axis. The end of the ladder is accelerating. Both axes are hidden from view.



$[-1, 15]$ by $[-1, 15]$

$$6. \frac{dy}{dt} = \frac{-4T}{\sqrt{10^2 - (2T)^2}}$$

7. $y'(3) \approx -4.24 \text{ ft/sec}^2$. The negative number means the ladder is falling.

8. Since $\lim_{t \rightarrow (13/3)^-} y'(t) = -\infty$, the speed of the top of the ladder is infinite as it hits the ground.

Quick Review 4.6

$$1. D = \sqrt{(7-0)^2 + (0-5)^2} = \sqrt{49+25} = \sqrt{74}$$

$$2. D = \sqrt{(b-0)^2 + (0-a)^2} = \sqrt{a^2 + b^2}$$

3. Use implicit differentiation.

$$\begin{aligned} \frac{d}{dx}(2xy + y^2) &= \frac{d}{dx}(x + y) \\ 2x \frac{dy}{dx} + 2y(1) + 2y \frac{dy}{dx} &= (1) + \frac{dy}{dx} \\ (2x + 2y - 1) \frac{dy}{dx} &= 1 - 2y \\ \frac{dy}{dx} &= \frac{1 - 2y}{2x + 2y - 1} \end{aligned}$$

4. Use implicit differentiation.

$$\begin{aligned} \frac{d}{dx}(x \sin y) &= \frac{d}{dx}(1 - xy) \\ (x)(\cos y) \frac{dy}{dx} + (\sin y)(1) &= -x \frac{dy}{dx} - y(1) \\ (x + x \cos y) \frac{dy}{dx} &= -y - \sin y \\ \frac{dy}{dx} &= \frac{-y - \sin y}{x + x \cos y} \\ \frac{dy}{dx} &= -\frac{y + \sin y}{x + x \cos y} \end{aligned}$$

5. Use implicit differentiation.

$$\begin{aligned} \frac{d}{dx} x^2 &= \frac{d}{dx} \tan y \\ 2x &= \sec^2 y \frac{dy}{dx} \\ \frac{dy}{dx} &= \frac{2x}{\sec^2 y} \\ \frac{dy}{dx} &= 2x \cos^2 y \end{aligned}$$

6. Use implicit differentiation.

$$\begin{aligned} \frac{d}{dx} \ln(x+y) &= \frac{d}{dx}(2x) \\ \frac{1}{x+y} \left(1 + \frac{dy}{dx} \right) &= 2 \\ 1 + \frac{dy}{dx} &= 2(x+y) \\ \frac{dy}{dx} &= 2x + 2y - 1 \end{aligned}$$

7. Using $A(-2, 1)$ we create the parametric equations $x = -2 + at$ and $y = 1 + bt$, which determine a line passing through A at $t = 0$. We determine a and b so that the line passes through $B(4, -3)$ at $t = 1$. Since $4 = -2 + a$, we have $a = 6$, and since $-3 = 1 + b$, we have $b = -4$. Thus, one parametrization for the line segment is $x = -2 + 6t$, $y = 1 - 4t$, $0 \leq t \leq 1$. (Other answers are possible.)

8. Using $A(0, -4)$, we create the parametric equations $x = 0 + at$ and $y = -4 + bt$, which determine a line passing through A at $t = 0$. We now determine a and b so that the line passes through $B(5, 0)$ at $t = 1$. Since $5 = 0 + a$, we have $a = 5$, and since $0 = -4 + b$, we have $b = 4$. Thus, one parametrization for the line segment is $x = 5t$, $y = -4 + 4t$, $0 \leq t \leq 1$. (Other answers are possible.)

9. One possible answer: $\frac{\pi}{2} \leq t \leq \frac{3\pi}{2}$

10. One possible answer: $\frac{3\pi}{2} \leq t \leq 2\pi$

Section 4.6 Exercises

1. Since $\frac{dA}{dt} = \frac{dA}{dr} \frac{dr}{dt}$, we have $\frac{dA}{dt} = 2\pi r \frac{dr}{dt}$,

3. (a) Since $\frac{dV}{dt} = \frac{dV}{dh} \frac{dh}{dt}$, we have $\frac{dV}{dt} = \pi r^2 \frac{dh}{dt}$.

(b) Since $\frac{dV}{dt} = \frac{dV}{dr} \frac{dr}{dt}$, we have $\frac{dV}{dt} = 2\pi r h \frac{dr}{dt}$.

(c)
$$\frac{dV}{dt} = \frac{d}{dt} \pi r^2 h = \pi \frac{d}{dt} (r^2 h)$$

$$\frac{dV}{dt} = \pi \left(r^2 \frac{dh}{dt} + h(2r) \frac{dr}{dt} \right)$$

$$\frac{dV}{dt} = \pi r^2 \frac{dh}{dt} + 2\pi r h \frac{dr}{dt}$$

5.
$$\frac{ds}{dt} = \frac{d}{dt} \sqrt{x^2 + y^2 + z^2}$$

$$\frac{ds}{dt} = \frac{1}{2\sqrt{x^2 + y^2 + z^2}} \frac{d}{dt} (x^2 + y^2 + z^2)$$

$$\frac{ds}{dt} = \frac{1}{2\sqrt{x^2 + y^2 + z^2}} \left(2x \frac{dx}{dt} + 2y \frac{dy}{dt} + 2z \frac{dz}{dt} \right)$$

$$\frac{ds}{dt} = \frac{x \frac{dx}{dt} + y \frac{dy}{dt} + z \frac{dz}{dt}}{\sqrt{x^2 + y^2 + z^2}}$$

7. (a) Since V is increasing at the rate of 1 volt/sec,

$$\frac{dV}{dt} = 1 \text{ volt/sec.}$$

- (b) Since I is decreasing at the rate of

$$\frac{1}{3} \text{ amp/sec, } \frac{dI}{dt} = -\frac{1}{3} \text{ amp/sec.}$$

- (c) Differentiating both sides of $V = IR$, we have

$$\frac{dV}{dt} = I \frac{dR}{dt} + R \frac{dI}{dt}.$$

- (d) Note that $V = IR$ gives $12 = 2R$, so $R = 6$ ohms. Now substitute the known values into the equation in (c).

$$1 = 2 \frac{dR}{dt} + 6 \left(-\frac{1}{3} \right)$$

$$3 = 2 \frac{dR}{dt}$$

$$\frac{dR}{dt} = \frac{3}{2} \text{ ohms/sec}$$

- R is changing at the rate of $\frac{3}{2}$ ohms/sec. Since this value is positive, R is increasing.

11. Continued

(b) $S = 4\pi r^2$

$$\frac{dS}{dt} = 8\pi r \frac{dr}{dt}$$

$$\frac{dS}{dt} = 8\pi(5)(1)$$

$$\frac{dS}{dt} = 40\pi \text{ ft}^2 / \text{min}$$

The surface area is increasing at the rate of 40π ft^2 / min

13. Step 1:

s = (diagonal) distance from antenna to airplane

x = horizontal distance from antenna to airplane

Step 2:

At the instant in question,

$$s = 10 \text{ mi and } \frac{ds}{dt} = 300 \text{ mph.}$$

Step 3:

We want to find $\frac{dx}{dt}$.

Step 4:

$$x^2 + 49 = s^2 \text{ or } x = \sqrt{s^2 - 49}$$

Step 5:


$$\frac{dx}{dt} = \frac{1}{2\sqrt{s^2 - 49}} \left(2s \frac{ds}{dt} \right) = \frac{s}{\sqrt{s^2 - 49}} \frac{ds}{dt}$$

Step 6:

$$\frac{dx}{dt} = \frac{10}{\sqrt{10^2 - 49}} (300) = \frac{3000}{\sqrt{51}} \text{ mph} \approx 420.08 \text{ mph}$$

The speed of the airplane is about 420.08 mph.

15. Step 1:



The cylinder shown represents the shape of the hole.

r = radius of cylinder

V = volume of cylinder

Step 2:

$$\text{At the instant in question, } \frac{dr}{dt} = \frac{0.001 \text{ in.}}{3 \text{ min}} = \frac{1}{3000} \text{ in./min}$$

and (since the diameter is 3.800 in.), $r = 1.900$ in.

Step 3:

We want to find $\frac{dV}{dt}$.

Step 4:

$$V = \pi r^2 (6) = 6\pi r^2$$

Step 5:

$$\frac{dV}{dt} = 12\pi r \frac{dr}{dt}$$

15. Continued


Step 6:

$$\frac{dV}{dt} = 12\pi(1.900)\left(\frac{1}{3000}\right) = \frac{19\pi}{2500} = 0.0076\pi$$

$$\approx 0.0239 \text{ in}^3/\text{min}.$$

The volume is increasing at the rate of approximately $0.0239 \text{ in}^3/\text{min}$.

17. Step 1:



r = radius of top surface of water

h = depth of water in reservoir

V = volume of water in reservoir

Step 2:

At the instant in question, $\frac{dV}{dt} = -50 \text{ m}^3/\text{min}$ and $h = 5 \text{ m}$.

Step 3:

We want to find $-\frac{dh}{dt}$ and $\frac{dr}{dt}$.

Step 4:

Note that $\frac{h}{r} = \frac{6}{45}$ by similar cones, so $r = 7.5h$.

$$\text{Then } V = \frac{1}{3}\pi r^2 h = \frac{1}{3}\pi(7.5h)^2 h = 18.75\pi h^3$$

Steps 5 and 6:

$$\text{(a) Since } V = 18.75\pi h^3, \frac{dV}{dt} = 56.25\pi h^2 \frac{dh}{dt}.$$

$$\text{Thus } -50 = 56.25\pi(5^2) \frac{dh}{dt}, \text{ and}$$

$$\text{so } \frac{dh}{dt} = -\frac{8}{225\pi} \text{ m/min} = -\frac{32}{9\pi} \text{ cm/min}.$$

The water level is falling by $\frac{32}{9\pi} \approx 1.13 \text{ cm/min}$.

(Since $\frac{dh}{dt} < 0$, the rate at which the water level is falling is positive.)

$$\text{(b) Since } r = 7.5h, \frac{dr}{dt} = 7.5 \frac{dh}{dt} = -\frac{80}{3\pi} \text{ cm/min. The rate of change of the radius of the water's surface is}$$

$$-\frac{80}{3\pi} \approx -8.49 \text{ cm/min}.$$

19. Step 1: x = distance from wall to base of ladder y = height of top of ladder A = area of triangle formed by the ladder, wall, and ground θ = angle between the ladder and the ground**Step 2:**At the instant in question, $x = 12$ ft and $\frac{dx}{dt} = 5$ ft/sec.**Step 3:**We want to find $-\frac{dy}{dt}$, $\frac{dA}{dt}$, and $\frac{d\theta}{dt}$.**Steps 4, 5, and 6:**

(a) $x^2 + y^2 = 169$

$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$$

To evaluate, note that, at the instant in question,

$$y = \sqrt{169 - x^2} = \sqrt{169 - 12^2} = 5.$$

$$\text{Then } 2(12)(5) + 2(5) \frac{dy}{dt} = 0$$

$$\frac{dy}{dt} = -12 \text{ ft/sec} \left(\text{or } -\frac{dy}{dt} = 12 \text{ ft/sec} \right)$$

The top of the ladder is sliding down the wall at the rate of 12 ft/sec. (Note that the *downward* rate of motion is positive.)

(b) $A = \frac{1}{2}xy$

$$\frac{dA}{dt} = \frac{1}{2} \left(x \frac{dy}{dt} + y \frac{dx}{dt} \right)$$

Using the results from step 2 and from part (a), we have

$$\frac{dA}{dt} = \frac{1}{2} [(12)(-12) + (5)(5)] = -\frac{119}{2} \text{ ft}^2/\text{sec.}$$

The area of the triangle is changing at the rate of $-59.5 \text{ ft}^2/\text{sec}$.

(c) $\tan \theta = \frac{y}{x}$

$$\sec^2 \theta \frac{d\theta}{dt} = \frac{x \frac{dy}{dt} - y \frac{dx}{dt}}{x^2}$$

Since $\tan \theta = \frac{5}{12}$, we have

$$\left(\text{for } 0 \leq \theta < \frac{\pi}{2} \right) \cos \theta = \frac{12}{13} \text{ and so } \sec^2 \theta = \frac{1}{\left(\frac{12}{13}\right)^2} = \frac{169}{144}.$$

Combining this result with the results from step 2 and

from part (a), we have $\frac{169}{144} \frac{d\theta}{dt} = \frac{(12)(-12) - (5)(5)}{12^2}$, so $\frac{d\theta}{dt} = -1$ radian/sec. The angle is changing at the rate of -1 radian/sec.

21. Step 1:

l = length of rope

x = horizontal distance from boat to dock

θ = angle between the rope and a vertical line

Step 2:

At the instant in question, $\frac{dl}{dt} = -2$ ft/sec and $l = 10$ ft.

Step 3:

We want to find the values of $-\frac{dx}{dt}$ and $\frac{d\theta}{dt}$.

Steps 4, 5, and 6:

$$(a) \quad x = \sqrt{l^2 - 36}$$

$$\frac{dx}{dt} = \frac{l}{\sqrt{l^2 - 36}} \frac{dl}{dt}$$

$$\frac{dx}{dt} = \frac{10}{\sqrt{10^2 - 36}} (-2) = -2.5 \text{ ft/sec}$$

The boat is approaching the dock at the rate of 2.5 ft/sec.

$$(b) \quad \theta = \cos^{-1} \frac{6}{l}$$

$$\frac{d\theta}{dt} = -\frac{1}{\sqrt{1 - \left(\frac{6}{l}\right)^2}} \left(-\frac{6}{l^2}\right) \frac{dl}{dt}$$

$$\frac{d\theta}{dt} = -\frac{1}{\sqrt{1 - 0.6^2}} \left(-\frac{6}{10^2}\right) (-2) = -\frac{3}{20} \text{ radian/sec}$$

The rate of change of angle θ is $-\frac{3}{20}$ radian/sec.

$$23. \quad \frac{dy}{dt} = \frac{dy}{dx} \frac{dx}{dt} = -10(1+x^2)^{-2} (2x) \frac{dx}{dt} = -\frac{20x}{(1+x^2)^2} \frac{dx}{dt}$$

Since $\frac{dx}{dt} = 3$ cm/sec, we have


$$\frac{dy}{dt} = -\frac{60x}{(1+x^2)^2} \text{ cm/sec.}$$

$$(a) \quad \frac{dy}{dt} = -\frac{60(-2)}{[1+(-2)^2]^2} = \frac{120}{5^2} = \frac{24}{5} \text{ cm/sec}$$

$$(b) \quad \frac{dy}{dt} = -\frac{60(0)}{(1+0^2)^2} = 0 \text{ cm/sec}$$

$$(c) \quad \frac{dy}{dt} = -\frac{60(20)}{(1+20^2)^2} \approx -0.00746 \text{ cm/sec}$$

25. Step 1:

 $x = x$ -coordinate of particle's location $y = y$ -coordinate of particle's location $\theta =$ angle of inclination of line joining the particle to the origin.

Step 2:

At the instant in question,

$$\frac{dx}{dt} = 10 \text{ m/sec and } x = 3 \text{ m.}$$

Step 3:

We want to find $\frac{d\theta}{dt}$.

Step 4:

Since $y = x^2$, we have $\tan \theta = \frac{y}{x} = \frac{x^2}{x} = x$ and so,for $x > 0$,

$$\theta = \tan^{-1} x.$$

Step 5:

$$\frac{d\theta}{dt} = \frac{1}{1+x^2} \frac{dx}{dt}$$

Step 6:

$$\frac{d\theta}{dt} = \frac{1}{1+3^2} (10) = 1 \text{ radian/sec}$$

The angle of inclination is increasing at the rate of 1 radian/sec.

27. Step 1:

 $r =$ radius of balls plus ice $S =$ surface area of ball plus ice $V =$ volume of ball plus ice

Step 2:

At the instant in question,

$$\frac{dV}{dt} = -8 \text{ mL/min} = -8 \text{ cm}^3/\text{min and } r = \frac{1}{2}(20) = 10 \text{ cm.}$$

Step 3:

We want to find $-\frac{dS}{dt}$.

Step 4:

We have $V = \frac{4}{3}\pi r^3$ and $S = 4\pi r^2$. These equations can becombined by noting that $r = \left(\frac{3V}{4\pi}\right)^{1/3}$, so $S = 4\pi \left(\frac{3V}{4\pi}\right)^{2/3}$

Step 5:


$$\frac{dS}{dt} = 4\pi \left(\frac{2}{3}\right) \left(\frac{3V}{4\pi}\right)^{-1/3} \left(\frac{3}{4\pi}\right) \frac{dV}{dt} = 2 \left(\frac{3V}{4\pi}\right)^{-1/3} \frac{dV}{dt}$$

Step 6:


$$\text{Note that } V = \frac{4}{3}\pi(10)^3 = \frac{4000\pi}{3}.$$

$$\frac{dS}{dt} = 2 \left(\frac{3}{4\pi} \cdot \frac{4000\pi}{3}\right)^{-1/3} (-8) = \frac{-16}{\sqrt[3]{1000}} = -1.6 \text{ cm}^2/\text{min}$$

Since $\frac{dS}{dt} < 0$, the rate of decrease is positive. The surface area is decreasing at the rate of $1.6 \text{ cm}^2/\text{min}$.



29. Step 1:



x = distance from streetlight base to man
 s = length of shadow

Step 2:

At the instant in question, $\frac{dx}{dt} = -5$ ft/sec and $x = 10$ ft.

Step 3:

We want to find $\frac{ds}{dt}$.

Step 4:

By similar triangles, $\frac{s}{6} = \frac{s+x}{16}$. This is equivalent to

$$16s = 6s + 6x, \text{ or } s = \frac{3}{5}x.$$

Step 5:

$$\frac{ds}{dt} = \frac{3}{5} \frac{dx}{dt}$$

Step 6:

$$\frac{ds}{dt} = \frac{3}{5}(-5) = -3 \text{ ft/sec}$$

The shadow length is changing at the rate of -3 ft/sec.



31. Step 1:

x = position of car ($x = 0$ when car is right in front of you)
 θ = camera angle. (We assume θ is negative until the car passes in front of you, and then positive.)

Step 2:

At the first instant in question, $x = 0$ ft and $\frac{dx}{dt} = 264$ ft/sec.

A half second later, $x = \frac{1}{2}(264) = 132$ ft and $\frac{dx}{dt} = 264$ ft/sec.

Step 3:

We want to find $\frac{d\theta}{dt}$ at each of the two instants.

Step 4:

$$\theta = \tan^{-1}\left(\frac{x}{132}\right)$$

Step 5:


$$\frac{d\theta}{dt} = \frac{1}{1 + \left(\frac{x}{132}\right)^2} \cdot \frac{1}{132} \frac{dx}{dt}$$

31. Continued

Step 6:

$$\text{When } x = 0: \frac{d\theta}{dt} = \frac{1}{1 + \left(\frac{0}{132}\right)^2} \left(\frac{1}{132}\right) (264) = 2 \text{ radians/sec}$$

$$\text{When } x = 132: \frac{d\theta}{dt} = \frac{1}{1 + \left(\frac{132}{132}\right)^2} \left(\frac{1}{132}\right) (264) = 1 \text{ radians/sec}$$



33. Step 1:

 s = shadow length θ = sun's angle of elevation

Step 2:

At the instant in question,

$$s = 60 \text{ ft and } \frac{d\theta}{dt} = 0.27^\circ / \text{min} = 0.0015\pi \text{ radian/min.}$$

Step 3:

We want to find $-\frac{ds}{dt}$.

Step 4:

$$\tan \theta = \frac{80}{s} \text{ or } s = 80 \cot \theta$$

Step 5:

$$\frac{ds}{dt} = -80 \csc^2 \theta \frac{d\theta}{dt}$$


Step 6:

Note that, at the moment in question, since \tan


$$\theta = \frac{80}{60} \text{ and } 0 < \theta < \frac{\pi}{2}, \text{ we have } \sin \theta = \frac{4}{5} \text{ and so}$$

$$\csc \theta = \frac{5}{4}.$$

$$\begin{aligned} \frac{ds}{dt} &= -80 \left(\frac{5}{4}\right)^2 (0.0015\pi) \\ &= -0.1875\pi \frac{\text{ft}}{\text{min}} \cdot \frac{12 \text{ in}}{1 \text{ ft}} \\ &= -2.25\pi \text{ in./min} \\ &\approx -7.1 \text{ in./min} \end{aligned}$$

Since $\frac{ds}{dt} < 0$, the rate at which the shadow length is*decreasing* is positive. The shadow length is decreasing at the rate of approximately 7.1 in./min.

35. Step 1:



a = distance from O to A
 b = distance from O to B
 c = distance from A to B

Step 2:

At the instant in question, $a = 5$ nautical miles, $b = 3$

nautical miles, $\frac{da}{dt} = 14$ knots, and $\frac{db}{dt} = 21$ knots.

Step 3:

We want to find $\frac{dc}{dt}$,

Step 4:

Law of Cosines : $c^2 = a^2 + b^2 - 2ab \cos 120^\circ$

$$c^2 = a^2 + b^2 + ab$$

Step 5:

$$2c \frac{dc}{dt} = 2a \frac{da}{dt} + 2b \frac{db}{dt} + a \frac{db}{dt} + b \frac{da}{dt}$$

Step 6:

Note that, at the instant in question,

$$c = \sqrt{a^2 + b^2 + ab} = \sqrt{(5)^2 + (3)^2 + (5)(3)} = \sqrt{49} = 7$$

$$2(7) \frac{dc}{dt} = 2(5)(14) + 2(3)(21) + (5)(21) + (3)(14)$$


$$14 \frac{dc}{dt} = 413$$

$$\frac{dc}{dt} = 29.5 \text{ knots}$$

The ships are moving apart at a rate of 29.5 knots.



37. False. Since $\frac{dA}{dt} = 2\pi r \frac{dr}{dt}$, the value of $\frac{dA}{dt}$ depends on r .



39. E. $sA = 6s^2$

$$dsA = 12s ds$$

$$12 = 12s ds$$


$$ds = \frac{1}{s}$$

$$V = s^3$$

$$dV = 3s^2 ds = 3s^2 \frac{1}{s}$$

$$24 = 3s$$

$$s = 8 \text{ in}$$



41. B. $v = \pi r^2 h$

$$sA = 2\pi r h$$

$$dv = \pi r^2 dh$$

$$dsA = 2\pi h dr$$


$$dv = dsA$$


$$\pi r^2 dh = 2\pi h dr$$

$$\frac{dh}{h} = 2 \frac{dr}{r^2}$$

$$\frac{2}{100} = 2 \frac{dr}{(1)^2}$$

$$dr = .01 \frac{\text{cm}}{\text{sec}}$$





43. (a) Note that the level of the coffee in the cone is not needed until part (b).

Step 1:

V_1 = volume of coffee in pot

y = depth of coffee in pot

Step 2:

$$\frac{dV_1}{dt} = 10 \text{ in}^3 / \text{min}$$

Step 3:

We want to find the value of $\frac{dy}{dt}$.

Step 4:

$$V_1 = 9\pi y$$

Step 5:

$$\frac{dV_1}{dt} = 9\pi \frac{dy}{dt}$$

Step 6:

$$10 = 9\pi \frac{dy}{dt}$$

$$\frac{dy}{dt} = \frac{10}{9\pi} \approx 0.354 \text{ in./min}$$

The level in the pot is increasing at the rate of approximately 0.354 in./min.

(b) Step 1:

V_2 = volume of coffee in filter

r = radius of surface of coffee in filter

h = depth of coffee in filter

Step 2:

At the instant in question, $\frac{dV_2}{dt} = -10 \text{ in}^3 / \text{min}$ and

$h = 5 \text{ in.}$

Step 3:

We want to find $-\frac{dh}{dt}$.

Step 4:

Note that $\frac{r}{h} = \frac{3}{6}$, so $r = \frac{h}{2}$.

$$\text{Then } V_2 = \frac{1}{3}\pi r^2 h = \frac{\pi h^3}{12}.$$

Step 5:

$$\frac{dV_2}{dt} = \frac{\pi h^2}{4} \frac{dh}{dt}$$

Step 6:


$$-10 = \frac{\pi(5)^2}{4} \frac{dh}{dt}$$

$$\frac{dh}{dt} = -\frac{8}{5\pi} \text{ in./min}$$

Note that $\frac{dh}{dt} < 0$, so the rate at which the level is

falling is positive. The level in the cone is falling at the

rate of $\frac{8}{5\pi} \approx 0.509 \text{ in./min.}$



45. (a) The point being plotted would correspond to a point on the edge of the wheel as the wheel turns.

(b) One possible answer is $\theta = 16\pi t$, where t is in seconds. (An arbitrary constant may be added to this expression, and we have assumed counterclockwise motion.)

45. Continued

(c) In general, assuming counterclockwise motion:

$$\frac{dx}{dt} = -2 \sin \theta \frac{d\theta}{dt} = -2(\sin \theta)(16\pi) = -32\pi \sin \theta$$

$$\frac{dy}{dt} = 2 \cos \theta \frac{d\theta}{dt} = 2(\cos \theta)(16\pi) = 32\pi \cos \theta$$

$$\text{At } \theta = \frac{\pi}{4}:$$

$$\frac{dx}{dt} = -32\pi \sin \frac{\pi}{4} = -16\pi(\sqrt{2}) \approx -71.086 \text{ ft/sec}$$

$$\frac{dy}{dt} = 32\pi \cos \frac{\pi}{4} = 16\pi(\sqrt{2}) \approx 71.086 \text{ ft/sec}$$

$$\text{At } \theta = \frac{\pi}{2}:$$

$$\frac{dx}{dt} = -32\pi \sin \frac{\pi}{2} = -32\pi \approx -100.531 \text{ ft/sec}$$

$$\frac{dy}{dt} = 32\pi \cos \frac{\pi}{2} = 0 \text{ ft/sec}$$

$$\text{At } \theta = \pi:$$

$$\frac{dx}{dt} = -32\pi \sin \pi = 0 \text{ ft/sec}$$

$$\frac{dy}{dt} = 32\pi \cos \pi = -32\pi \approx -100.531 \text{ ft/sec}$$

$$\begin{aligned} \text{(b)} \quad \frac{dy}{dt} &= \frac{d}{dt}(uv) = u \frac{dv}{dt} + v \frac{du}{dt} \\ &= u(0.03v) + v(-0.02u) \\ &= 0.01uv \\ &= 0.01y \end{aligned}$$

The total production is increasing at the rate of 1% per year.

Quick Quiz Sections 4.4–4.6

$$1. \text{ B. } x_{n+1} = x_n - \frac{f(x)}{f'(x)}$$

$$f(x) = x^3 + 2x - 1$$

$$f'(x) = 3x^2 + 2$$

$$x_2 = 1 - \frac{(1)^3 + 2(1) - 1}{3(1)^2 + 2} = \frac{3}{5}$$

$$x_3 = \frac{3}{5} - \frac{\left(\frac{3}{5}\right)^3 + 2\left(\frac{3}{5}\right) - 1}{3\left(\frac{3}{5}\right)^2 + 2} = 0.465$$

$$2. \text{ B. } z^2 = x^2 + y^2$$

$$z = \sqrt{4^2 + 3^2} = 5$$

$$2z \frac{dz}{dt} = 2x \frac{dx}{dt} + 2y \frac{dy}{dt}$$

$$5 = 4 \left(3 \frac{dy}{dt} \right) + 3 \frac{dy}{dt}$$

$$\frac{dy}{dt} = \frac{1}{3}$$

$$\frac{dx}{dt} = 3 \frac{dy}{dt} = 3 \left(\frac{1}{3} \right) = 1$$

$$3. \text{ A. } x(t) = 70$$

$$y(t) = 60t$$

$$z(t) = ((60t)^2 + 70^2)^{1/2}$$

$$\frac{dz}{dt} = \frac{1}{2} (3600t^2 + 4900)^{-1/2} (7200t)$$

$$\frac{dz}{dt} = \frac{7200(4)}{2(3600(4)^2 + 4900)^{1/2}}$$

$$\frac{dz}{dt} = 57.6$$

$$4. \text{ (a) } f(x) = \sqrt{x}$$

$$x = 25$$

$$f'(25) = \frac{1}{2} (25)^{-1/2} = \frac{1}{10}$$

$$\sqrt{26} = 5 + \frac{1}{10} (26 - 25) = 5.1$$

$$\begin{aligned} 47. \text{ (a)} \quad \frac{dy}{dt} &= \frac{d}{dt}(uv) = u \frac{dv}{dt} + v \frac{du}{dt} \\ &= u(0.05v) + v(0.04u) \\ &= 0.09uv \\ &= 0.09y \end{aligned}$$

Since $\frac{dy}{dt} = 0.09y$, the rate of growth of total production is 9% per year.

4. Continued

(b) $x_{n+1} = x_n - \frac{f(x)}{f'(x)}, f(x) = x^2 - 26 = 0$

$x_2 = 5 - \frac{(5)^2 - 26}{2(5)} = 5.1$

(c) $f(x) = \sqrt[3]{x}$
 $x = 3$

$f'(27) = \frac{1}{3}(27)^{-2/3} = \frac{1}{27}$

$\sqrt{26} = 3 + \frac{1}{27}(26 - 27)$

$\sqrt{26} = 2.963$

Chapter 4 Review (pp. 256–260)

1. $y = x\sqrt{2-x}$

$y' = x \left(\frac{1}{2\sqrt{2-x}} \right) (-1) + (\sqrt{2-x})(1)$
 $= \frac{-x + 2(2-x)}{2\sqrt{2-x}}$
 $= \frac{4-3x}{2\sqrt{2-x}}$

The first derivative has a zero at $\frac{4}{3}$.

Critical point value: $x = \frac{4}{3} \quad y = \frac{4\sqrt{6}}{9} \approx 1.09$

Endpoint values: $x = -2 \quad y = -4$
 $x = 2 \quad y = 0$

The global maximum value is $\frac{4\sqrt{6}}{9}$ at $x = \frac{4}{3}$, and the global minimum value is -4 at $x = -2$.

2. Since y is a cubic function with a positive leading coefficient, we have $\lim_{x \rightarrow -\infty} y = -\infty$ and $\lim_{x \rightarrow \infty} y = \infty$. There are no global extrema.

3. $y' = (x^2)(e^{1/x^2})(-2x^{-3}) + (e^{1/x^2})(2x)$


$= 2e^{1/x^2} \left(-\frac{1}{x} + x \right)$
 $= \frac{2e^{1/x^2}(x-1)(x+1)}{x}$

| | | | | |
|-----------------|------------|--------------|-------------|------------|
| Intervals | $x < -1$ | $-1 < x < 0$ | $0 < x < 1$ | $x > 1$ |
| Sign of y' | - | + | - | + |
| Behavior of y | Decreasing | Increasing | Decreasing | Increasing |

$y'' = \frac{d}{dx} [2e^{1/x^2}(-x^{-1} + x)]$
 $= (2e^{1/x^2})(x^{-2} + 1) + (-x^{-1} + x)(2e^{1/x^2})(-2x^{-3})$
 $= (2e^{1/x^2})(x^{-2} + 1) + 2x^{-4} - 2x^{-2}$
 $= \frac{2e^{1/x^2}(x^4 - x^2 + 2)}{x^4}$
 $= \frac{2e^{1/x^2}[(x^2 - 0.5)^2 + 1.75]}{x^4}$

The second derivative is always positive (where defined), so the function is concave up for all $x \neq 0$.

Graphical support:



- (a) $[-1, 0)$ and $[1, \infty)$
- (b) $(-\infty, -1]$ and $(0, 1]$
- (c) $(-\infty, 0)$ and $(0, \infty)$
- (d) None
- (e) Local (and absolute) minima at $(1, e)$ and $(-1, e)$
- (f) None

4. Note that the domain of the function is $[-2, 2]$.

$y' = x \left(\frac{1}{2\sqrt{4-x^2}} \right) (-2x) + (\sqrt{4-x^2})(1)$
 $= \frac{-x^2 + (4-x^2)}{\sqrt{4-x^2}}$
 $= \frac{4-2x^2}{\sqrt{4-x^2}}$

| | | | |
|-----------------|----------------------|----------------------------|--------------------|
| Intervals | $-2 < x < -\sqrt{2}$ | $-\sqrt{2} < x < \sqrt{2}$ | $\sqrt{2} < x < 2$ |
| Sign of y' | - | + | - |
| Behavior of y | Decreasing | Increasing | Decreasing |


$y'' = \frac{(\sqrt{4-x^2})(-4x) - (4-2x^2) \left(\frac{1}{2\sqrt{4-x^2}} \right) (-2x)}{4-x^2}$
 $= \frac{2x(x^2-6)}{(4-x^2)^{3/2}}$

Note that the values $x = \pm\sqrt{6}$ are not zeros of y'' because they fall outside of the domain.

| | | |
|-----------------|--------------|--------------|
| Intervals | $-2 < x < 0$ | $0 < x < 2$ |
| Sign of y'' | + | - |
| Behavior of y | Concave up | Concave down |

4. Continued

Graphical support:



$[-2.35, 2.35]$ by $[-3.5, 3.5]$

- (a) $[-\sqrt{2}, \sqrt{2}]$
- (b) $[-2, -\sqrt{2}]$ and $[\sqrt{2}, 2]$
- (c) $(-2, 0)$
- (d) $(0, 2)$
- (e) Local maxima: $(-2, 0)$, $(\sqrt{2}, 2)$

Local minima: $(2, 0)$, $(-\sqrt{2}, -2)$

Note that the extrema at $x = \pm\sqrt{2}$ are also absolute extrema.

- (f) $(0, 0)$

5. $y' = 1 - 2x - 4x^3$


Using grapher techniques, the zero of y' is $x \approx 0.385$.

| Intervals | $x < 0.385$ | $0.385 < x$ |
|-----------------|-------------|-------------|
| Sign of y' | + | - |
| Behavior of y | Increasing | Decreasing |

$y'' = -2 - 12x^2 = -2(1 + 6x^2)$

The second derivative is always negative so the function is concave down for all x .

Graphical support:



$[-4, 4]$ by $[-4, 2]$

- (a) Approximately $(-\infty, 0.385]$
- (b) Approximately $[0.385, \infty)$
- (c) None
- (d) $(-\infty, \infty)$
- (e) Local (and absolute) maximum at $\approx (0.385, 1.215)$
- (f) None


6. $y' = e^{x-1} - 1$

| Intervals | $x < 1$ | $1 < x$ |
|-----------------|------------|------------|
| Sign of y' | - | + |
| Behavior of y | Decreasing | Increasing |

$y'' = e^{x-1}$

The second derivative is always positive, so the function is concave up for all x .

Graphical support:



$[-4, 4]$ by $[-2, 4]$

- (a) $[1, \infty)$
- (b) $(-\infty, 1]$
- (c) $(-\infty, \infty)$
- (d) None
- (e) Local (and absolute) minimum at $(1, 0)$
- (f) None

7. Note that the domain is $(-1, 1)$.

$y = (1 - x^2)^{-1/4}$

$y' = -\frac{1}{4}(1 - x^2)^{-5/4}(-2x) = \frac{x}{2(1 - x^2)^{5/4}}$

| Intervals | $-1 < x < 0$ | $0 < x < 1$ |
|-----------------|--------------|-------------|
| Sign of y' | - | + |
| Behavior of y | Decreasing | Increasing |


$$y'' = \frac{2(1 - x^2)^{5/4}(1 - (x)(2)\left(\frac{5}{4}\right)(1 - x^2)^{1/4}(-2x))}{4(1 - x^2)^{5/2}}$$

$$= \frac{(1 - x^2)^{1/4}[2 - 2x^2 + 5x^2]}{4(1 - x^2)^{5/2}}$$

$$= \frac{3x^2 + 2}{4(1 - x^2)^{9/4}}$$

The second derivative is always positive, so the function is concave up on its domain $(-1, 1)$.

Graphical support:



$[-1.3, 1.3]$ by $[-1, 3]$

- (a) $[0, 1)$
- (b) $(-1, 0]$
- (c) $(-1, 1)$
- (d) None
- (e) Local minimum at $(0, 1)$
- (f) None

$$8. y' = \frac{(x^3 - 1)(1) - (x)(3x^2)}{(x^3 - 1)^2} = \frac{2x^3 + 1}{(x^3 - 1)^2}$$

| | | | |
|-----------------|-----------------|---------------------|------------|
| Intervals | $x < -2^{-1/3}$ | $-2^{-1/3} < x < 1$ | $1 < x$ |
| Sign of y' | + | - | - |
| Behavior of y | Increasing | Decreasing | Decreasing |


$$y'' = -\frac{(x^3 - 1)^2(6x^2) - (2x^3 + 1)(2)(x^3 - 1)(3x^2)}{(x^3 - 1)^4}$$

$$= -\frac{(x^3 - 1)(6x^2) - (2x^3 + 1)(6x^2)}{(x^3 - 1)^3}$$

$$= \frac{6x^2(x^3 + 2)}{(x^3 - 1)^3}$$

| | | | | |
|-----------------|----------------|--------------------|--------------|------------|
| Intervals | $x < -2^{1/3}$ | $-2^{1/3} < x < 0$ | $0 < x < 1$ | $1 < x$ |
| Sign of y'' | + | - | - | + |
| Behavior of y | Concave up | Concave down | Concave down | Concave up |

Graphical support:



$[-4.7, 4.7]$ by $[-3.1, 3.1]$

- (a) $(-\infty, -2^{-1/3}] \approx (-\infty, -0.794]$
- (b) $[-2^{-1/3}, 1) \approx [-0.794, 1)$ and $(1, \infty)$
- (c) $(-\infty, -2^{-1/3}) \approx (-\infty, -1.260)$ and $(1, \infty)$
- (d) $(-2^{-1/3}, 1) \approx (-1.260, 1)$
- (e) Local minimum at $(-2^{-1/3}, \frac{2}{3} \cdot 2^{-1/3}) \approx (-0.794, 0.529)$
- (f) $(-2^{1/3}, \frac{1}{3} \cdot 2^{1/3}) \approx (-1.260, 0.420)$

9. Note that the domain is $[-1, 1]$.

$$y' = -\frac{1}{\sqrt{1-x^2}}$$


Since y' is negative on $(-1, 1)$ and y is continuous, y is decreasing on its domain $[-1, 1]$.

$$y'' = \frac{d}{dx} [-(1-x^2)^{-1/2}]$$

$$= \frac{1}{2}(1-x^2)^{-3/2}(-2x) = -\frac{x}{(1-x^2)^{3/2}}$$

| | | |
|-----------------|--------------|--------------|
| Intervals | $-1 < x < 0$ | $0 < x < 1$ |
| Sign of y'' | + | - |
| Behavior of y | Concave up | Concave down |


Graphical support:




$[-1.175, 1.175]$ by $[-\frac{\pi}{4}, \frac{5\pi}{4}]$

- (a) None
- (b) $[-1, 1]$
- (c) $(-1, 0)$
- (d) $(0, 1)$
- (e) Local (and absolute) maximum at $(-1, \pi)$; local (and absolute) minimum at $(1, 0)$
- (f) $(0, \frac{\pi}{2})$


10. This problem can be solved graphically by using NDER to obtain the graphs shown below.



$[-4, 4]$ by $[-1, 0.3]$



$[-4, 4]$ by $[-0.4, 0.6]$



$[-4, 4]$ by $[-0.7, 0.8]$

An alternative approach using a combination of algebraic and graphical techniques follows. Note that the denominator of y is always positive because it is equivalent to $(x+1)^2 + 2$.

$$y' = \frac{(x^2 + 2x + 3)(1) - (x)(2x + 2)}{(x^2 + 2x + 3)^2}$$

$$= \frac{-x^2 + 3}{(x^2 + 2x + 3)^2}$$

| | | | |
|-----------------|-----------------|----------------------------|----------------|
| Intervals | $x < -\sqrt{3}$ | $-\sqrt{3} < x < \sqrt{3}$ | $\sqrt{3} < x$ |
| Sign of y' | - | + | - |
| Behavior of y | Decreasing | Increasing | Decreasing |

$$y'' = \frac{(x^2 + 2x + 3)^2(-2x) - (-x^2 + 3)(2)(x^2 + 2x + 3)(2x + 2)}{(x^2 + 2x + 3)^4}$$

$$= \frac{(x^2 + 2x + 3)(-2x) - 2(2x + 2)(-x^2 + 3)}{(x^2 + 2x + 3)^3}$$

$$= \frac{2x^3 - 18x - 12}{(x^2 + 2x + 3)^3}$$

10. Continued

Using graphing techniques, the zeros of $2x^3 - 18x - 12$ (and hence of y'') are at $x \approx -2.584$, $x \approx -0.706$, and $x \approx 3.290$.

| | | | | |
|-----------------|---------------------|--------------------|-------------------|-------------------|
| Intervals | $(-\infty, -2.584)$ | $(-2.584, -0.706)$ | $(-0.706, 3.290)$ | $(3.290, \infty)$ |
| Sign of y'' | - | + | - | + |
| Behavior of y | Concave down | Concave up | Concave down | Concave up |

- (a) $[-\sqrt{3}, \sqrt{3}]$
- (b) $(-\infty, -\sqrt{3})$ and $[\sqrt{3}, \infty)$
- (c) Approximately $(-2.584, -0.706)$ and $(3.290, \infty)$
- (d) Approximately $(-\infty, -2.584)$ and $(-0.706, 3.290)$
- (e) Local maximum at $\left(\sqrt{3}, \frac{\sqrt{3}-1}{4}\right) \approx (1.732, 0.183)$;
local minimum at $\left(-\sqrt{3}, \frac{-\sqrt{3}-1}{4}\right) \approx (-1.732, -0.683)$
- (f) $\approx (-2.584, -0.573)$, $(-0.706, -0.338)$, and $(3.290, 0.161)$

11. For $x > 0$, $y' = \frac{d}{dx} \ln x = \frac{1}{x}$
For $x < 0$: $y' = \frac{d}{dx} \ln(-x) = \frac{1}{-x}(-1) = \frac{1}{x}$


Thus $y' = \frac{1}{x}$ for all x in the domain.

| | | |
|-----------------|------------|------------|
| Intervals | $(-2, 0)$ | $(0, 2)$ |
| Sign of y' | - | + |
| Behavior of y | Decreasing | Increasing |

$y'' = -x^{-2}$

The second derivative always negative, so the function is concave down on each open interval of its domain.

Graphical support:



$[-2.35, 2.35]$ by $[-3, 1.5]$

- (a) $(0, 2]$
- (b) $[-2, 0)$
- (c) None
- (d) $(-2, 0)$ and $(0, 2)$

(e) Local (and absolute) maxima at $(-2, \ln 2)$ and $(2, \ln 2)$

(f) None

12. $y' = 3 \cos 3x - 4 \sin 4x$

Using graphing techniques, the zeros of y' in the domain

$0 \leq x \leq 2\pi$ are $x \approx 0.176$, $x \approx 0.994$, $x = \frac{\pi}{2} \approx 1.57$,
 $x \approx 2.148$, and $x \approx 2.965$, $x \approx 3.834$, $x = \frac{3\pi}{2}$, $x \approx 5.591$

| | | | | | |
|-----------------|-----------------|---------------------|-----------------------------|-----------------------------|---------------------|
| Intervals | $0 < x < 0.176$ | $0.176 < x < 0.994$ | $0.994 < x < \frac{\pi}{2}$ | $\frac{\pi}{2} < x < 2.148$ | $2.148 < x < 2.965$ |
| Sign of y' | + | - | + | - | + |
| Behavior of y | Increasing | Decreasing | Increasing | Decreasing | Increasing |

| | | | | |
|-----------------|---------------------|------------------------------|------------------------------|--------------------|
| Intervals | $2.965 < x < 3.834$ | $3.834 < x < \frac{3\pi}{2}$ | $\frac{3\pi}{2} < x < 5.591$ | $5.591 < x < 2\pi$ |
| Sign of y' | - | + | - | + |
| Behavior of y | Decreasing | Increasing | Decreasing | Increasing |

$y'' = -9 \sin 3x - 16 \cos 4x$


Using graphing techniques, the zeros of y'' in the domain

$0 \leq x \leq 2\pi$ are $x \approx 0.542$, $x \approx 1.266$, $x \approx 1.876$,
 $x \approx 2.600$, $x \approx 3.425$, $x \approx 4.281$, $x \approx 5.144$ and $x \approx 6.000$.

| | | | | | |
|-----------------|-----------------|---------------------|---------------------|---------------------|---------------------|
| Intervals | $0 < x < 0.542$ | $0.542 < x < 1.266$ | $1.266 < x < 1.876$ | $1.876 < x < 2.600$ | $2.600 < x < 3.425$ |
| Sign of y'' | - | + | - | + | - |
| Behavior of y | Concave down | Concave up | Concave down | Concave up | Concave down |

| | | | | |
|-----------------|---------------------|---------------------|---------------------|-------------------|
| Intervals | $3.425 < x < 4.281$ | $4.281 < x < 5.144$ | $5.144 < x < 6.000$ | $6.00 < x < 2\pi$ |
| Sign of y'' | + | - | + | - |
| Behavior of y | Concave up | Concave down | Concave up | Concave down |

Graphical support:



$\left[-\frac{\pi}{4}, \frac{9\pi}{4}\right]$ by $[-2.5, 2.5]$

12. Continued

(a) Approximately $[0, 0.176]$,

$$\left[0.994, \frac{\pi}{2}\right], [2.148, 2.965], \left[3.834, \frac{3\pi}{2}\right], \text{ and } [5.591, 2\pi]$$

(b) Approximately $[0.176, 0.994]$,

$$\left[\frac{\pi}{2}, 2.148\right], [2.965, 3.834], \text{ and } \left[\frac{3\pi}{2}, 5.591\right]$$

(c) Approximately $(0.542, 1.266)$, $(1.876, 2.600)$, $(3.425, 4.281)$, and $(5.144, 6.000)$

(d) Approximately $(0, 0.542)$, $(1.266, 1.876)$, $(2.600, 3.425)$, $(4.281, 5.144)$, and $(6.000, 2\pi)$

(e) Local maxima at $\approx (0.176, 1.266)$, $\left(\frac{\pi}{2}, 0\right)$

and $(2.965, 1.266)$, $\left(\frac{3\pi}{2}, 2\right)$, and $(2\pi, 1)$;

local minima at $\approx (0, 1)$, $(0.994, -0.513)$, $(2.148, -0.513)$, $(3.834, -1.806)$, and $(5.591, -1.806)$

Note that the local extrema at $x \approx 3.834$, $x = \frac{3\pi}{2}$,

and $x \approx 5.591$ are also extrema.

(f) $\approx (0.542, 0.437)$, $(1.266, -0.267)$, $(1.876, -0.267)$, $(2.600, 0.437)$, $(3.425, -0.329)$, $(4.281, 0.120)$, $(5.144, 0.120)$, and $(6.000, -0.329)$


13. $y' = \begin{cases} -e^{-x}, & x < 0 \\ 4 - 3x^2, & x > 0 \end{cases}$

| Intervals | $x < 0$ | $0 < x < \frac{2}{\sqrt{3}}$ | $\frac{2}{\sqrt{3}} < x$ |
|-----------------|------------|------------------------------|--------------------------|
| Sign of y' | - | + | - |
| Behavior of y | Decreasing | Increasing | Decreasing |

$$y'' = \begin{cases} e^{-x}, & x > 0 \\ -6x, & x < 0 \end{cases}$$

| Intervals | $x < 0$ | $0 < x$ |
|-----------------|------------|--------------|
| Sign of y'' | + | - |
| Behavior of y | Concave up | Concave down |

Graphical support:



(a) $\left[0, \frac{2}{\sqrt{3}}\right]$

(b) $(-\infty, 0]$ and $\left[\frac{2}{\sqrt{3}}, \infty\right)$

(c) $(-\infty, 0)$

(d) $(0, \infty)$

(e) Local maximum at $\left(\frac{2}{\sqrt{3}}, \frac{16}{3\sqrt{3}}\right) \approx (1.155, 3.079)$

(f) None. Note that there is no point of inflection at $x = 0$ because the derivative is undefined and no tangent line exists at this point.

14. $y' = -5x^4 + 7x^2 + 10x + 4$

Using graphing techniques, the zeros of y' are $x \approx -0.578$ and $x \approx -1.692$.


| Intervals | $x < -0.578$ | $-0.578 < x < 1.692$ | $1.692 < x$ |
|-----------------|--------------|----------------------|-------------|
| Sign of y' | - | + | - |
| Behavior of y | Decreasing | Increasing | Decreasing |

$$y'' = -20x^3 + 14x + 10$$

Using graphing techniques, the zeros of y'' is $x \approx 1.079$.

| Intervals | $x < 1.079$ | $1.079 < x$ |
|-----------------|-------------|--------------|
| Sign of y'' | + | - |
| Behavior of y | Concave up | Concave down |

Graphical support:



(a) Approximately $[-0.578, 1.692]$

(b) Approximately $(-\infty, -0.578]$ and $[1.692, \infty)$

(c) Approximately $(-\infty, 1.079)$

(d) Approximately $(1.079, \infty)$

(e) Local maximum at $\approx (1.692, 20.517)$; local minimum at $\approx (-0.578, 0.972)$

(f) $\approx (1.079, 13.601)$

15. $y = 2x^{4/5} - x^{9/5}$


$$y' = \frac{8}{5}x^{-1/5} - \frac{9}{5}x^{4/5} = \frac{8-9x}{5\sqrt[5]{x}}$$

| | | | |
|-----------------|------------|-----------------------|-------------------|
| Intervals | $x < 0$ | $0 < x < \frac{8}{9}$ | $\frac{8}{9} < x$ |
| Sign of y' | - | + | - |
| Behavior of y | Decreasing | Increasing | Decreasing |

$$y'' = -\frac{8}{25}x^{-6/5} - \frac{36}{25}x^{-1/5} = \frac{4(2+9x)}{25x^{6/5}}$$

| | | | |
|-----------------|--------------------|------------------------|--------------|
| Intervals | $x < -\frac{2}{9}$ | $-\frac{2}{9} < x < 0$ | $0 < x$ |
| Sign of y'' | + | - | - |
| Behavior of y | Concave up | Concave down | Concave down |

Graphical support:



[-4, 4] by [-3, 3]

(a) $\left[0, \frac{8}{9}\right]$

(b) $(-\infty, 0]$ and $\left[\frac{8}{9}, \infty\right)$

(c) $\left(-\infty, -\frac{2}{9}\right)$

(d) $\left(-\frac{2}{9}, 0\right)$ and $(0, \infty)$


(e) Local maximum

at $\left(\frac{8}{9}, \frac{10}{9} \cdot \left(\frac{8}{9}\right)^{4/5}\right) \approx (0.889, 1.011)$; local minimum

at $(0, 0)$

(f) $\left(-\frac{2}{9}, \frac{20}{9} \cdot \left(-\frac{2}{9}\right)^{4/5}\right) \approx \left(-\frac{2}{9}, 0.667\right)$

16. We use a combination of analytic and grapher techniques to solve this problem. Depending on the viewing windows chosen, graphs obtained using NDER may exhibit strange behavior near $x = 2$ because, for example, NDER $(y, 2) \approx 5,000,000$ while y' is actually undefined at $x = 2$. The graph of $y = \frac{5-4x+4x^2-x^3}{x-2}$ is shown below.




[-5.875, 5.875] by [-50, 30]

$$y' = \frac{(x-2)(-4+8x-3x^2) - (5-4x+4x^2-x^3)(1)}{(x-2)^2}$$

$$= \frac{-2x^3+10x^2-16x+3}{(x-2)^2}$$

The graph of y' is shown below.



[-5.875, 5.875] by [-50, 30]

The zero of y' is $x \approx 0.215$.


| | | | |
|-----------------|-------------|-----------------|------------|
| Intervals | $x < 0.215$ | $0.215 < x < 2$ | $2 < x$ |
| Sign of y' | + | - | - |
| Behavior of y | Increasing | Decreasing | Decreasing |

$$y'' = \frac{(x-2)^2(-6x^2+20x-16) - (-2x^3+10x^2-16x+3)(2)(x-2)}{(x-2)^4}$$

$$= \frac{(x-2)(-6x^2+20x-16) - 2(-2x^3+10x^2-16x+3)}{(x-2)^3}$$

$$= \frac{-2(x^3-6x^2+12x-13)}{(x-2)^3}$$

The graph of y'' is shown below.



[-5.875, 5.875] by [-20, 20]

The zero of $x^3 - 6x^2 + 12x - 13$ (and hence of y'') is $x \approx 3.710$.

| | | | |
|-----------------|--------------|-----------------|--------------|
| Intervals | $x < 2$ | $2 < x < 3.710$ | $3.710 < x$ |
| Sign of y'' | - | + | - |
| Behavior of y | Concave down | Concave up | Concave down |

16. Continued

- (a) Approximately $(-\infty, 0.215]$
- (b) Approximately $[0.215, 2)$ and $(2, \infty)$
- (c) Approximately $(2, 3.710)$
- (d) $(-\infty, 2)$ and approximately $(3.710, \infty)$
- (e) Local maximum at $\approx (0.215, -2.417)$
- (f) $\approx (3.710, -3.420)$

17. $y' = 6(x+1)(x-2)^2$

| | | | |
|-----------------|------------|--------------|------------|
| Intervals | $x < -1$ | $-1 < x < 2$ | $2 < x$ |
| Sign of y' | - | + | + |
| Behavior of y | Decreasing | Increasing | Increasing |

$$y'' = 6(x+1)(2)(x-2) + 6(x-2)^2(1)$$

$$= 6(x-2)[(2x+2) + (x-2)]$$

$$= 18x(x-2)$$

| | | | |
|-----------------|------------|--------------|------------|
| Intervals | $x < 0$ | $0 < x < 2$ | $2 < x$ |
| Sign of y'' | + | - | + |
| Behavior of y | Concave up | Concave down | Concave up |

- (a) There are no local maxima.
- (b) There is a local (and absolute) minimum at $x = -1$.
- (c) There are points of inflection at $x = 0$ and at $x = 2$.

18. $y' = 6(x+1)(x-2)$

| | | | |
|-----------------|------------|--------------|------------|
| Intervals | $x < -1$ | $-1 < x < 2$ | $2 < x$ |
| Sign of y' | + | - | + |
| Behavior of y | Increasing | Decreasing | Increasing |

$$y'' = \frac{d}{dx} 6(x^2 - x - 2) = 6(2x - 1)$$

| | | |
|-----------------|-------------------|-------------------|
| Intervals | $x < \frac{1}{2}$ | $\frac{1}{2} < x$ |
| Sign of y'' | - | + |
| Behavior of y | Concave down | Concave up |

- (a) There is a local maximum at $x = -1$.
- (b) There is a local maximum at $x = 2$.
- (c) There is a point of inflection at $x = \frac{1}{2}$.

19. Since $\frac{d}{dx} \left(-\frac{1}{4}x^{-4} - e^{-x} \right) = x^{-5} + e^{-x}$,

$$f(x) = -\frac{1}{4}x^{-4} - e^{-x} + C.$$

20. Since $\frac{d}{dx} \sec x = \sec x \tan x$, $f(x) = \sec x + C$.

21. Since $\frac{d}{dx} \left(2 \ln x + \frac{1}{3}x^3 + x \right) = \frac{2}{x} + x^2 + 1$,

$$f(x) = 2 \ln x + \frac{1}{3}x^3 + x + C.$$

22. Since $\frac{d}{dx} \left(\frac{2}{3}x^{3/2} + 2x^{1/2} \right) = \sqrt{x} + \frac{1}{\sqrt{x}}$,

$$f(x) = \frac{2}{3}x^{3/2} + 2x^{1/2} + C.$$

23. $f(x) = -\cos x + \sin x + C$

$$f(\pi) = 3$$

$$1 + 0 + C = 3$$

$$C = 2$$

$$f(x) = -\cos x + \sin x + 2$$

24. $f(x) = \frac{3}{4}x^{4/3} + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + C$

$$f(1) = 0$$

$$\frac{3}{4} + \frac{1}{3} + \frac{1}{2} + 1 + C = 0$$

$$C = -\frac{31}{12}$$

$$f(x) = \frac{3}{4}x^{4/3} + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x - \frac{31}{12}$$

25. $v(t) = s'(t) = 9.8t + 5$

$$s(t) = 4.9t^2 + 5t + C$$

$$s(0) = 10$$

$$C = 10$$

$$s(t) = 4.9t^2 + 5t + 10$$

26. $a(t) = v'(t) = 32$

$$v(t) = 32t + C_1$$

$$v(0) = 20$$

$$C_1 = 20$$

$$v(t) = s'(t) = 32t + 20$$

$$s(t) = 16t^2 + 20t + C_2$$

$$s(0) = 5$$

$$C_2 = 5$$

$$s(t) = 16t^2 + 20t + 5$$

27. $f(x) = \tan x$
 $f'(x) = \sec^2 x$

$$\begin{aligned} L(x) &= f\left(-\frac{\pi}{4}\right) + f'\left(-\frac{\pi}{4}\right)\left[x - \left(-\frac{\pi}{4}\right)\right] \\ &= \tan\left(-\frac{\pi}{4}\right) + \sec^2\left(-\frac{\pi}{4}\right)\left(x + \frac{\pi}{4}\right) \\ &= -1 + 2\left(x + \frac{\pi}{4}\right) \\ &= 2x + \frac{\pi}{2} - 1 \end{aligned}$$

28. $f(x) = \sec x$
 $f'(x) = \sec x \tan x$

$$\begin{aligned} L(x) &= f\left(\frac{\pi}{4}\right) + f'\left(\frac{\pi}{4}\right)\left(x - \frac{\pi}{4}\right) \\ &= \sec\left(\frac{\pi}{4}\right) + \sec\left(\frac{\pi}{4}\right)\tan\left(\frac{\pi}{4}\right)\left(x - \frac{\pi}{4}\right) \\ &= \sqrt{2} + \sqrt{2}(1)\left(x - \frac{\pi}{4}\right) \\ &= \sqrt{2}x - \frac{\pi\sqrt{2}}{4} + \sqrt{2} \end{aligned}$$

29. $f(x) = \frac{1}{1 + \tan x}$
 $f'(x) = -(1 + \tan x)^{-2}(\sec^2 x)$
 $= -\frac{1}{\cos^2 x(1 + \tan x)^2}$
 $= -\frac{1}{(\cos x + \sin x)^2}$
 $L(x) = f(0) + f'(0)(x - 0)$
 $= 1 - 1(x - 0)$
 $= -x + 1$

30. $f(x) = e^x + \sin x$
 $f'(x) = e^x + \cos x$
 $L(x) = f(0) + f'(0)(x - 0)$
 $= 1 + 2(x - 0)$
 $= 2x + 1$

31. The global minimum value of $\frac{1}{2}$ occurs at $x = 2$.

32. (a) The values of y' and y'' are both negative where the graph is decreasing and concave down, at T .

(b) The value of y' is negative and the value of y'' is positive where the graph is decreasing and concave up, at P .


33. (a) The function is increasing on the interval $(0, 2]$.

(b) The function is decreasing on the interval $[-3, 0)$.

(c) The local extreme values occur only at the endpoints of the domain. A local maximum value of 1 occurs at $x = -13$, and a local maximum value of 3 occurs at $x = 2$.

34. The 24th day


35.



36. (a) We know that f is decreasing on $[0, 1]$ and increasing on $[1, 3]$, the absolute minimum value occurs at $x = 1$ and the absolute maximum value occurs at an endpoint. Since $f(0) = 0$, $f(1) = -2$, and $f(3) = 3$, the absolute minimum value is -2 at $x = 1$ and the absolute maximum value is 3 at $x = 3$.

(b) The concavity of the graph does not change. There are no points of inflection.

(c)



37. (a) $f(x)$ is continuous on $[0.5, 3]$ and differentiable on $(0.5, 3)$.

(b) $f'(x) = (x)\left(\frac{1}{x}\right) + (\ln x)(1) = 1 + \ln x$

Using $a = 0.5$ and $b = 3$, we solve as follows.

$$f'(c) = \frac{f(3) - f(0.5)}{3 - 0.5}$$

$$1 + \ln c = \frac{3 \ln 3 - 0.5 \ln 0.5}{2.5}$$

$$\ln c = \frac{\ln\left(\frac{3^3}{0.5^{0.5}}\right)}{2.5} - 1$$

$$\ln c = 0.4 \ln(27\sqrt{2}) - 1$$

$$c = e^{-1}(27\sqrt{2})^{0.4}$$

$$c = e^{-1}\sqrt[5]{1458} \approx 1.579$$

(c) The slope of the line is

$$m = \frac{f(b) - f(a)}{b - a} = 0.4 \ln(27\sqrt{2}) - 0.2 \ln 1458, \text{ and the line}$$

passes through $(3, 3 \ln 3)$. Its equation is

$$y = 0.2(\ln 1458)(x - 3) + 3 \ln 3, \text{ or approximately}$$

$$y = 1.457x - 1.075.$$

37. Continued

- (d) The slope of the line is
- $m = 0.2 \ln 1458$
- , and the line passes through

$$(c, f(c)) = (e^{-1} \sqrt[5]{1458}, e^{-1} \sqrt[5]{1458}(-1 + 0.2 \ln 1458)) \\ \approx (1.579, 0.722).$$

Its equation is

$$y = 0.2(\ln 1458)(x - c) + f(c), \\ y = 0.2 \ln 1458(x - e^{-1} \sqrt[5]{1458}) \\ + e^{-1} \sqrt[5]{1458}(-1 + 0.2 \ln 1458),$$

$$y = 0.2(\ln 1458)x - e^{-1} \sqrt[5]{1458}, \\ \text{or approximately } y = 1.457x - 1.579.$$

38. (a) $v(t) = s'(t) = 4 - 6t - 3t^2$

(b) $a(t) = v'(t) = -6 - 6t$

- (c) The particle starts at position 3 moving in the positive direction, but decelerating. At approximately
- $t = 0.528$
- , it reaches position 4.128 and changes direction, beginning to move in the negative direction. After that, it continues to accelerate while moving in the negative direction.

39. (a) $L(x) = f(0) + f'(0)(x - 0) \\ = -1 + 0(x - 0) = -1$

(b) $f(0.1) \approx L(0.1) = -1$


- (c) Greater than the approximation in (b), since
- $f'(x)$
- is actually positive over the interval
- $(0, 0.1)$
- and the estimate is based on the derivative being 0.

40. (a) Since $\frac{dy}{dx} = (x^2)(-e^{-x}) + (e^{-x})(2x) + (2x - x^2)e^{-x}$,

$$dy = (2x - x^2)e^{-x} dx.$$

(b) $dy = [2(1) - (1)^2](e^{-1})(0.01) \\ = 0.01e^{-1} \\ \approx 0.00368$

41. (a) With some rounding, $y = \frac{1633001.59}{1 + 17.471e^{-0.06378t}}$




(c) $y = \frac{1633001.59}{1 + 17.471e^{-0.06378(80)}} + 829,210 = 2,305,337$

- (d) Using the Second Derivative, we find the maximum rate of growth about 1885. We find a point of inflection here, which shows the beginning of a decline in the rate of growth.

(e) $y = \frac{1633001.59}{1 + 17.471e^{-0.06378(\infty)}} \approx 2,462,000$, which is the approximate maximum population.

- (f) There are many possible causes. Advances in transportation began drawing the population southward after 1920, and Tennessee was well-situated geographically to become a crossroads of river, railroad, and automobile routes. By the year 2000 there had been numerous other demographic changes. It should be pointed out that the census years in the data (1850–1910) include the years of the Civil War and Reconstruction, so the regression is based on unusual data.

42. $f(x) = 2 \cos x - \sqrt{1+x}$
 $f'(x) = -2 \sin x - \frac{1}{2\sqrt{1+x}}$
 $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$
 $= x_n - \frac{2 \cos x_n - \sqrt{1+x_n}}{-2 \sin x_n - \frac{1}{2\sqrt{1+x_n}}}$

The graph of $y = f(x)$ shows that $f(x) = 0$ has one solution, near $x = 1$.

$$x_1 = 1 \\ x_2 \approx 0.8361848 \\ x_3 \approx 0.8283814 \\ x_4 \approx 0.8283608 \\ x_5 \approx 0.8283608$$

Solution: $x \approx 0.828361$

43. Let
- t
- represent time in seconds, where the rocket lifts off at
- $t = 0$
- . Since
- $a(t) = v'(t) = 20$
- , m/sec
- ²
- and
- $v(0) = 0$
- m/sec, we have
- $v(t) = 20t$
- , and so
- $v(60) = 1200$
- m/sec. The speed after 1 minute (60 seconds) will be 1200 m/sec.

44. Let t represent time in seconds, where the rock is blasted upward at $t = 0$. Since $a(t) = v'(t) = -3.72 \text{ m/sec}^2$ and $v(0) = 93 \text{ m/sec}$, we have $v(t) = -3.72t + 93$. Since $s'(t) = -3.72t + 93$ and $s(0) = 0$, we have $s(t) = -1.86t^2 + 93t$. Solving $v(t) = 0$, we find that the rock attains its maximum height at $t = 25$ sec and its height at that time is $s(25) = 1162.5 \text{ m}$.


45. Note that $s = 100 - 2r$ and the sector area is given by

$$A = \pi r^2 \left(\frac{s}{2\pi r} \right) = \frac{1}{2}rs = \frac{1}{2}r(100 - 2r) = 50r - r^2. \text{ To find}$$

the domain of $A(r) = 50r - r^2$, note that $r > 0$ and

$$0 < s < 2\pi r, \text{ which gives } 12.1 \approx \frac{50}{\pi + 1} < r < 50. \text{ Since}$$

$A'(r) = 50 - 2r$, the critical point occurs at $r = 25$. This value is in the domain and corresponds to the maximum area because $A''(r) = -2$, which is negative for all r . The greatest area is attained when $r = 25$ ft and $s = 50$ ft.



For $0 < x < \sqrt{27}$, the triangle with vertices at $(0, 0)$ and $(\pm x, 27 - x^2)$ has an area given by

$$A(x) = \frac{1}{2}(2x)(27 - x^2) = 27x - x^3. \text{ Since}$$

$A' = 27 - 3x^2 = 3(3 - x)(3 + x)$ and $A'' = -6x$, the critical point in the interval $(0, \sqrt{27})$ occurs at $x = 3$ and corresponds to the maximum area because $A''(x)$ is negative in this interval. The largest possible area is $A(3) = 54$ square units.

47. If the dimensions are x ft by x ft by h ft, then the total amount of steel used is $x^2 + 4xh \text{ ft}^2$. Therefore,

$$x^2 + 4xh = 108 \text{ and so } h = \frac{108 - x^2}{4x}. \text{ The volume is given}$$

$$\text{by } V(x) = x^2h = \frac{108x - x^3}{4} = 27x - 0.25x^3. \text{ Then}$$

$$V'(x) = 27 - 0.75x^2 = 0.75(6 + x)(6 - x) \text{ and}$$

$V''(x) = -1.5x$. The critical point occurs at $x = 6$, and it corresponds to the maximum volume because $V''(x) < 0$

for $x > 0$. The corresponding height is $\frac{108 - 6^2}{4(6)} = 3$ ft. The

base measures 6 ft by 6 ft, and the height is 3 ft.

48. If the dimensions are x ft by x ft by h ft, then we have

$$x^2h = 32 \text{ and so } h = \frac{32}{x^2}. \text{ Neglecting the quarter-inch}$$

thickness of the steel, the area of the steel used is

$$A(x) = x^2 + 4xh = x^2 + \frac{128}{x}. \text{ We can minimize the weight}$$

of the vat by minimizing this quantity. Now

$$A'(x) = 2x - 128x^{-2} = \frac{2}{x^2}(x^3 - 4^3) \text{ and}$$

$A''(x) = 2 + 256x^{-3}$. The critical point occurs at $x = 4$ and corresponds to the minimum possible area because

$$A''(x) > 0 \text{ for } x > 0. \text{ The corresponding height is } \frac{32}{4^2} = 2 \text{ ft.}$$

The base should measure 4 ft by 4 ft, and the height should be 2 ft.

49. We have $r^2 + \left(\frac{h}{2}\right)^2 = 3$, so $r^2 = 3 - \frac{h^2}{4}$. We wish to

minimize the cylinder's volume

$$V = \pi r^2 h = \pi \left(3 - \frac{h^2}{4} \right) h = 3\pi h - \frac{\pi h^3}{4} \text{ for } 0 < h < 2\sqrt{3}.$$

$$\text{Since } \frac{dV}{dh} = 3\pi - \frac{3\pi h^2}{4} = \frac{3\pi}{4}(2 + h)(2 - h) \text{ and}$$

$$\frac{d^2V}{dh^2} = -\frac{3\pi h}{2}, \text{ the critical point occurs at } h = 2 \text{ and it}$$

corresponds to the maximum value because $\frac{d^2V}{dh^2} < 0$ for

$h > 0$. The corresponding value of r is $\sqrt{3 - \frac{2^2}{4}} = \sqrt{2}$. The

largest possible cylinder has height 2 and radius $\sqrt{2}$.

50. Note that, from similar cones, $\frac{r}{6} = \frac{12 - h}{12}$, so $h = 12 - 2r$.

The volume of the smaller cone is given by

$$V = \frac{1}{3}\pi r^2 h = \frac{1}{3}\pi r^2(12 - 2r) = 4\pi r^2 - \frac{2\pi}{3}r^3 \text{ for } 0 < r < 6.$$


Then $\frac{dV}{dr} = 8\pi r - 2\pi r^2 = 2\pi r(4 - r)$, so the critical point

occurs at $r = 4$. This critical point corresponds to the

maximum volume because $\frac{dV}{dr} > 0$ for $0 < r < 4$ and


$\frac{dV}{dr} < 0$ for $4 < r < 6$. The smaller cone has the largest possible value when $r = 4$ ft and $h = 4$ ft.

51.



(a) $V(x) = x(15 - 2x)(5 - x)$

(b, c) Domain: $0 < x < 5$



The maximum volume is approximately 66.019 and it occurs when $x \approx 1.962$ in.

(d) Note that $V(x) = 2x^3 - 25x^2 + 75x$,

so $V'(x) = 6x^2 - 50x + 75$.

Solving $V'(x) = 0$, we have


$$x = \frac{50 \pm \sqrt{(-50)^2 - 4(6)(75)}}{2(6)} = \frac{50 \pm \sqrt{700}}{12}$$

$$= \frac{50 \pm 10\sqrt{7}}{12} = \frac{25 \pm 5\sqrt{7}}{6}$$

These solutions are approximately $x \approx 1.962$ and $x = 6.371$, so the critical point in the appropriate domain occurs at

$$x = \frac{25 - 5\sqrt{7}}{6}$$

52.



For $0 < x < \frac{5\pi}{3}$, the area of the rectangle is given by

$$A(x) = (2x)(8 \cos 0.3x) = 16x \cos 0.3x$$

Then $A'(x) = 16x(-0.3 \sin 0.3x) + 16(\cos 0.3x)(1)$
 $= 16(\cos 0.3x - 0.3x \sin 0.3x)$

Solving $A'(x) = 0$ graphically, we find that the critical point occurs at $x \approx 2.868$ and the corresponding area is approximately 29.925 square units.

53. The cost (in thousands of dollars) is given by

$$C(x) = 40x + 30(20 - y) = 40x + 600 - 30\sqrt{x^2 - 144}$$

Then $C'(x) = 40 - \frac{30}{2\sqrt{x^2 - 144}}(2x) = 40 - \frac{30x}{\sqrt{x^2 - 144}}$

Solving $C'(x) = 0$, we have:

$$\frac{30x}{\sqrt{x^2 - 144}} = 40$$

$$3x = 4\sqrt{x^2 - 144}$$

$$9x^2 = 16x^2 - 2304$$

$$2304 = 7x^2$$

Choose the positive solution:

$$x = +\frac{48}{\sqrt{7}} \approx 18.142 \text{ mi}$$

$$y = \sqrt{x^2 - 12^2} = \frac{36}{\sqrt{7}} \approx 13.607 \text{ mi}$$

54. The length of the track is given by $2x + 2\pi r$, so we have $2x + 2\pi r = 400$ and therefore $x = 200 - \pi r$. Then the area of the rectangle is

$$A(r) = 2rx$$

$$= 2r(200 - \pi r)$$

$$= 400r - 2\pi r^2, \text{ for } 0 < r < \frac{200}{\pi}$$

Therefore, $A'(r) = 400 - 4\pi r$ and $A''(r) = -4\pi$, so the

critical point occurs at $r = \frac{100}{\pi}$ m and this point

corresponds to the maximum rectangle area because $A''(r) < 0$ for all r .

The corresponding value of x is

$$x = 200 - \pi \left(\frac{100}{\pi} \right) = 100 \text{ m}$$

The rectangle will have the largest possible area when

$$x = 100 \text{ m and } r = \frac{100}{\pi}$$

55. Assume the profit is k dollars per hundred grade B tires and $2k$ dollars per hundred grade A tires.

Then the profit is given by

$$P(x) = 2kx + k \cdot \frac{40 - 10x}{5 - x}$$

$$= 2k \cdot \frac{(20 - 5x) + x(5 - x)}{5 - x}$$

$$= 2k \cdot \frac{20 - x^2}{5 - x}$$

$$P'(x) = 2k \cdot \frac{(5 - x)(-2x) - (20 - x^2)(-1)}{(5 - x)^2}$$

$$= 2k \cdot \frac{x^2 - 10x + 20}{(5 - x)^2}$$

55. Continued

The solutions of $P'(x) = 0$ are

$$x = \frac{10 \pm \sqrt{(-10)^2 - 4(1)(20)}}{2(1)} = 5 \pm \sqrt{5}, \text{ so the solution in the}$$

appropriate domain is $x = 5 - \sqrt{5} \approx 2.76$.

Check the profit for the critical point and endpoints:

Critical point: $x \approx 2.76$ $P(x) \approx 11.06k$

End points: $x = 0$ $P(x) = 8k$

$x = 4$ $P(x) = 8k$


The highest profit is obtained when $x \approx 2.76$ and $y \approx 5.53$, which corresponds to 276 grade A tires and 553 grade B tires.

56. (a) The distance between the particles is $|f(t)|$ where

$$f(t) = -\cos t + \cos\left(t + \frac{\pi}{4}\right). \text{ Then}$$

$$f'(t) = \sin t - \sin\left(t + \frac{\pi}{4}\right)$$

Solving $f'(t) = 0$ graphically, we obtain $t \approx 1.178$, $t \approx 4.230$, and so on.



$[0, 2\pi]$ by $[-2, 2]$

Alternatively, $f'(t) = 0$ may be solved analytically as follows.

$$\begin{aligned} f'(t) &= \sin\left[\left(t + \frac{\pi}{8}\right) - \frac{\pi}{8}\right] - \sin\left[\left(t + \frac{\pi}{8}\right) + \frac{\pi}{8}\right] \\ &= \left[\sin\left(t + \frac{\pi}{8}\right)\cos\frac{\pi}{8} - \cos\left(t + \frac{\pi}{8}\right)\sin\frac{\pi}{8}\right] \\ &\quad - \left[\sin\left(t + \frac{\pi}{8}\right)\cos\frac{\pi}{8} + \cos\left(t + \frac{\pi}{8}\right)\sin\frac{\pi}{8}\right] \\ &= -2\sin\frac{\pi}{8}\cos\left(t + \frac{\pi}{8}\right), \end{aligned}$$


so the critical points occur when

$$\cos\left(t + \frac{\pi}{8}\right) = 0, \text{ or } t = \frac{3\pi}{8} + k\pi. \text{ At each of these values,}$$

$f(t) = \pm 2\cos\frac{3\pi}{8} \approx \pm 0.765$ units, so the maximum distance between the particles is 0.765 units.

(b) Solving $\cos t = \cos\left(t + \frac{\pi}{4}\right)$ graphically, we obtain

$t \approx 2.749$, $t \approx 5.890$, and so on.



$[0, 2\pi]$ by $[-2, 2]$

Alternatively, this problem may be solved analytically as follows.

$$\begin{aligned} \cos t &= \cos\left(t + \frac{\pi}{4}\right) \\ \cos\left[\left(t + \frac{\pi}{8}\right) - \frac{\pi}{8}\right] &= \cos\left[\left(t + \frac{\pi}{8}\right) + \frac{\pi}{8}\right] \\ \cos\left(t + \frac{\pi}{8}\right)\cos\frac{\pi}{8} + \sin\left(t + \frac{\pi}{8}\right)\sin\frac{\pi}{8} &= \cos\left(t + \frac{\pi}{8}\right)\cos\frac{\pi}{8} \\ &\quad - \sin\left(t + \frac{\pi}{8}\right)\sin\frac{\pi}{8} \\ 2\sin\left(t + \frac{\pi}{8}\right)\sin\frac{\pi}{8} &= 0 \\ \sin\left(t + \frac{\pi}{8}\right) &= 0 \\ t &= \frac{7\pi}{8} + k\pi \end{aligned}$$


The particles collide when $t = \frac{7\pi}{8} \approx 2.749$ (plus multiples of π if they keep going.)

57. The dimensions will be x in. by $10 - 2x$ in. by $16 - 2x$ in., so $V(x) = x(10 - 2x)(16 - 2x) = 4x^3 - 52x^2 + 160x$ for $0 < x < 5$.

Then $V'(x) = 12x^2 - 104x + 160 = 4(x - 2)(3x - 20)$, so the critical point in the correct domain is $x = 2$.

This critical point corresponds to the maximum possible volume because $V'(x) > 0$ for $0 < x < 2$ and $V'(x) < 0$ for $2 < x < 5$. The box of largest volume has a height of 2 in. and a base measuring 6 in. by 12 in., and its volume is 144 in^3 .

Graphical support:



$[0, 5]$ by $[-40, 160]$

58. Step 1:

 r = radius of circle A = area of circle

Step 2:

At the instant in question, $\frac{dr}{dt} = -\frac{2}{\pi}$ m/sec and $r = 10$ m.

Step 3:

We want to find $\frac{dA}{dt}$.

Step 4:

$$A = \pi r^2$$

Step 5:

$$\frac{dA}{dt} = 2\pi r \frac{dr}{dt}$$

Step 6:

$$\frac{dA}{dt} = 2\pi(10)\left(-\frac{2}{\pi}\right) = -40$$

The area is changing at the rate of -40 m²/sec.

59. Step 1:

 x = x -coordinate of particle y = y -coordinate of particle D = distance from origin to particle

Step 2:

At the instant in question, $x = 5$ m, $y = 12$ m,

$$\frac{dx}{dt} = -1 \text{ m/sec, and } \frac{dy}{dt} = -5 \text{ m/sec.}$$

Step 3:

We want to find $\frac{dD}{dt}$.

Step 4:

$$D = \sqrt{x^2 + y^2}$$

Step 5:

$$\frac{dD}{dt} = \frac{1}{2\sqrt{x^2 + y^2}} \left(2x \frac{dx}{dt} + 2y \frac{dy}{dt} \right) = \frac{x \frac{dx}{dt} + y \frac{dy}{dt}}{\sqrt{x^2 + y^2}}$$

Step 6:

$$\frac{dD}{dt} = \frac{(5)(-1) + (12)(-5)}{\sqrt{5^2 + 12^2}} = -5 \text{ m/sec}$$

Since $\frac{dD}{dt}$ is negative, the particle is *approaching* the origin at the *positive* rate of 5 m/sec.

60. Step 1:

 x = edge of length of cube V = volume of cube

Step 2:

At the instant in question,

$$\frac{dV}{dt} = 1200 \text{ cm}^3/\text{min and } x = 20 \text{ cm.}$$

Step 3:

We want to find $\frac{dx}{dt}$.

Step 4:

$$V = x^3$$

Step 5:

$$\frac{dV}{dt} = 3x^2 \frac{dx}{dt}$$

Step 6:

$$1200 = 3(20)^2 \frac{dx}{dt}$$

$$\frac{dx}{dt} = 1 \text{ cm/min}$$

The edge length is increasing at the rate of 1 cm/min.

61. Step 1:

 x = x -coordinate of point y = y -coordinate of point D = distance from origin to point

Step 2:

At the instant in question, $x = 3$ and $\frac{dD}{dt} = 11$ units per sec.

Step 3:

We want to find $\frac{dx}{dt}$.

Step 4:

Since $D^2 = x^2 + y^2$ and $y = x^{3/2}$, we have

$$D = \sqrt{x^2 + x^3} \text{ for } x \geq 0.$$

Step 5:

$$\begin{aligned} \frac{dD}{dt} &= \frac{1}{2\sqrt{x^2 + x^3}} (2x + 3x^2) \frac{dx}{dt} \\ &= \frac{2x + 3x^2}{2x\sqrt{1+x}} \frac{dx}{dt} = \frac{3x+2}{2\sqrt{1+x}} \frac{dx}{dt} \end{aligned}$$

61. Continued

Step 6:

$$11 = \frac{3(3) + 2}{2\sqrt{4}} \frac{dx}{dt}$$

$$\frac{dx}{dt} = 4 \text{ units per sec}$$

62. (a) Since $\frac{h}{r} = \frac{10}{4}$, we may write $h = \frac{5r}{2}$ or $r = \frac{2h}{5}$.

(b) Step 1:

h = depth of water in tank
 r = radius of surface of water
 V = volume of water in tank

Step 2:

At the instant in question,

$$\frac{dV}{dt} = -5 \text{ ft}^3/\text{min} \text{ and } h = 6 \text{ ft.}$$

Step 3:

We want to find $-\frac{dh}{dt}$.

Step 4:

$$V = \frac{1}{3}\pi r^2 h = \frac{4}{75}\pi h^3$$

Step 5:

$$\frac{dV}{dt} = \frac{4}{25}\pi h^2 \frac{dh}{dt}$$

Step 6:

$$-5 = \frac{4}{25}\pi(6)^2 \frac{dh}{dt}$$

$$\frac{dh}{dt} = -\frac{125}{144\pi} \approx -0.276 \text{ ft/min}$$

Since $\frac{dh}{dt}$ is negative, the water level is *dropping* at the positive rate of ≈ 0.276 ft/min.

63. Step 1:

r = radius of outer layer of cable on the spool
 θ = clockwise angle turned by spool
 s = length of cable that has been unwound

Step 2:

At the instant in question, $\frac{ds}{dt} = 6$ ft/sec and $r = 1.2$ ft

Step 3:

We want to find $\frac{d\theta}{dt}$.

Step 4:

$$s = r\theta$$

Step 5:

Since r is essentially constant, $\frac{ds}{dt} = r \frac{d\theta}{dt}$

Step 6:

$$6 = 1.2 \frac{d\theta}{dt}$$

$$\frac{d\theta}{dt} = 5 \text{ radians/sec}$$

The spool is turning at the rate of 5 radians per second.

64. $a(t) = v'(t) = -g = -32 \text{ ft/sec}^2$

Since $v(0) = 32 \text{ ft/sec}$, $v(t) = s'(t) = -32t + 32$.

Since $s(0) = -17 \text{ ft}$, $s(t) = -16t^2 + 32t - 17$.

The shovelful of dirt reaches its maximum height when $v(t) = 0$, at $t = 1$ sec. Since $s(1) = -1$, the shovelful of dirt is still below ground level at this time. There was not enough speed to get the dirt out of the hole. Duck!

65. We have $V = \frac{1}{3}\pi r^2 h$, so $\frac{dV}{dr} = \frac{2}{3}\pi r h$ and $dV = \frac{2}{3}\pi r h dr$.

When the radius changes from a to $a + dr$, the volume

change is approximately $dV = \frac{2}{3}\pi a h dr$.

66. (a) Let x = edge of length of cube and S = surface area of

cube. Then $S = 6x^2$, which means $\frac{dS}{dx} = 12x$ and

$dS = 12x dx$. We want $|dS| \leq 0.02S$, which gives

$|12x dx| \leq 0.02(6x^2)$ or $|dx| \leq 0.01x$. The edge should be measured with an error of no more than 1%.

(b) Let V = volume of cube. Then $V = x^3$, which means

$\frac{dV}{dx} = 3x^2$ and $dV = 3x^2 dx$. We have $|dx| \leq 0.01x$,

which means $|3x^2 dx| \leq 3x^2(0.01x) = 0.03V$,

so $|dV| \leq 0.03V$. The volume calculation will be accurate to within approximately 3% of the correct volume.

67. Let C = circumference, r = radius, S = surface area, and V = volume.

(a) Since $C = 2\pi r$, we have $\frac{dC}{dr} = 2\pi$ and so $dC = 2\pi dr$.

Therefore, $\left|\frac{dC}{C}\right| = \left|\frac{2\pi dr}{2\pi r}\right| = \left|\frac{dr}{r}\right| < \frac{0.4 \text{ cm}}{10 \text{ cm}} = 0.04$. The calculated radius will be within approximately 4% of the correct radius.

(b) Since $S = 4\pi r^2$, we have $\frac{dS}{dr} = 8\pi r$ and so

$dS = 8\pi r dr$. Therefore, $\left|\frac{dS}{S}\right| = \left|\frac{8\pi r dr}{4\pi r^2}\right| = \left|\frac{2 dr}{r}\right| \leq 2(0.04) = 0.08$. The

calculated surface area will be within approximately 8% of the correct surface area.

(c) Since $V = \frac{4}{3}\pi r^3$, we have $\frac{dV}{dr} = 4\pi r^2$ and so

$dV = 4\pi r^2 dr$. Therefore

$\left|\frac{dV}{V}\right| = \left|\frac{4\pi r^2 dr}{\frac{4}{3}\pi r^3}\right| = \left|\frac{3 dr}{r}\right| \leq 3(0.04) = 0.12$.

The calculated volume will be within approximately 12% of the correct volume.

68. By similar triangles, we have $\frac{a}{6} = \frac{a+20}{h}$, which gives

$ah = 6a + 120$, or $h = 6 + 120a^{-1}$. The height of the lamp post is approximately $6 + 120(15)^{-1} = 14$ ft. The estimated error in measuring a was

$|da| \leq 1$ in. $= \frac{1}{12}$ ft. Since $\frac{dh}{da} = -120a^{-2}$, we have

$|dh| = |-120a^{-2} da| \leq 120(15)^{-2} \left(\frac{1}{12}\right) = \frac{2}{45}$ ft, so the

estimated possible error is $\pm \frac{2}{45}$ ft or $\pm \frac{8}{15}$ in.


69. $\frac{dy}{dx} = 2 \sin x \cos x - 3$. Since $\sin x$ and $\cos x$ are both between 1 and -1 , the value of $2 \sin x \cos x$ is never greater than 2. Therefore, $\frac{dy}{dx} \leq 2 - 3 = -1$ for all values of x .

Since $\frac{dy}{dx}$ is always negative, the function decreases on every interval.

70. (a) f has a relative maximum at $x = -2$. This is where $f'(x) = 0$, causing f' to go from positive to negative.

(b) f has a relative minimum at $x = 0$. This is where $f'(x) = 0$, causing f' to go from negative to positive.

(c) The graph of f is concave up on $(-1, 1)$ and on $(2, 3)$. These are the intervals on which the derivatives of f are increasing.



71. (a) $A = \pi r^2$

$$\frac{dA}{dt} = 2\pi r dr$$

$$\frac{dA}{dt} = 2\pi(2)\left(\frac{1}{3}\right) = \frac{4}{3}\pi \frac{\text{in.}^2}{\text{sec}}$$

(b) $dA = dV$

$$\frac{4}{3}\pi = \frac{1}{3}\pi r^2 dh$$

$$\frac{4}{3}\pi = \frac{1}{3}\pi(2)^2 dh$$

$$\frac{dh}{dt} = 1 \frac{\text{in.}}{\text{sec}}$$

(c) $\frac{dA}{dh} = \frac{\frac{4}{3}\pi}{1} = \frac{4}{3}\pi \frac{\text{in.}^2}{\text{in.}}$

72. (a) $2a + 4b = 60$

$$b = 15 - 2a$$

$$V = \pi a^2 b = \pi a^2 (15 - 2a)$$

$$\frac{dV}{da} = 30\pi a - \frac{3\pi a^2}{2}$$


$$30\pi a = \frac{3\pi a^2}{2}$$

$$a = 20$$

$$2(20) + 4b = 60$$

$$b = 5$$

(b) The sign graph for the derivative $\frac{dV}{da} = \frac{3\pi a}{2}(20 - a)$ on the interval $(0, 30)$ is as follows:



By the First Derivative Test, there is a maximum at $x = 20$.